

<https://doi.org/10.3176/chem.1979.3.06>

УДК 547.872/874
541.182.644

*Tiina LINNAS, Aili KOGERMAN,
H. HEINLO, O. KIRRET*

REACTION OF HYDROXYALKYL METHACRYLATE GELS WITH CHLORINE DERIVATES OF TRIAZINE

Of the carriers for affinity chromatography, the synthetic macroporous polymer hydrogels based on copolymers of glycol methacrylate with glycol dimethacrylate are of great interest. These gels exhibit a remarkable chemical and mechanical stability. Affinants can be bound to hydroxyl groups of gel with cyanogen bromide by using a method analogical to that of binding the affinant to Sepharose [1].

The method of binding biologically active molecules to HMG (hydroxyalkyl methacrylate gels) with cyanogen bromide has several disadvantages, toxicity of the reagents used being one of them. We have found that the binding of biologically active molecules to the —OH groups of HMG may be regarded possible when less toxic reagents TCT (2,4,6-trichlorotriazine) and DCT (2-amino-4,6-dichlorotriazine) are used.

The objective of this work was to investigate the binding of trypsin to the above-mentioned gels by the aid of chlorine derivates of triazine and to demonstrate the use of immobilized trypsin in the isolation of trypsin antibody from rabbit serum immunized with a suspension of tissue antigens of pancreas extract [2].

Material and methods

Enzymes. Bovine trypsin (EC 3.4.4.4) «B» was obtained from the Olaine Works (U. S. S. R.). Esterolytic activity of trypsin was 23 (μ moles·min) \cdot mg and immobilized trypsin was assayed with BAEE (N-benzoyl-L-arginine ethylester) at 25°C with the pH-stat method.

Chemicals. Two hydroxyalkyl methacrylate gels, produced and marked under the trade mark Spheron from LACHEMA Company (Brno, Czechoslovakia), were used: basic Spheron P-1000 (particle size 0.063 to 0.100 mm) and Spheron C-1000 (particle size 0.040 to 0.060 mm) containing *p*-nitrophenylester groups 0.095 m·eqv/g and —COOH—groups 0.2 m·eqv/g. BAEE·HCl and Hammarsten casein were from Reanal (Hungary). TCT (analytical grade) used was a product of the Yerevan Chemical Works and DCT was synthesized in our laboratory by the method described by G. Kay and M. D. Lilly [3].

Activating HMG with chlorine derivates of triazine. In order to activate HMG with TCT or DCT, the following method was used: 1 g of HMG was activated with 20 ml of 1 to 3 N NaOH solution for 15 min. Then the solution of NaOH was sucked off and 50 ml of a 2% solution of TCT in a mixture of xyloldioxane (1:1) or 50 ml of DCT in acetone or benzene were added to the product. The reaction was carried out at room temperature. The product obtained was sucked off, washed with a mixture of water—dioxane—acetic acid (1:1:1), with water and acetone, and dried under vacuum.

The reaction product was assayed (H, C, N) on a Hewlett—Packard automatic analyzer. The chlorine content was determined by potentiometric titration.

Binding trypsin to Spheron. The binding of trypsin to Spheron P-1000 activated with chlorine derivates of triazine was carried out in 0.2 M H_3BO_3/Na_2CO_3 buffer at 23°. The binding reaction was broken off by rapid acidification of samples with pH adjusted to 2.5—3.0. After that the samples were washed with tris-buffer.

The binding of trypsin to Spheron C-1000 proceeded as follows: a certain amount of Spheron was stirred in a ten-fold excess of water and stored in a refrigerator for swelling for 2 to 3 h. The swollen gel was filtered, transferred to a phosphate buffer (pH 7.0) containing about 200 mg of trypsin per 1 g of dry gel, and shaken for 60 min at 20°.

Unnecessary valency groups were washed off with a 1 M solution of ethanolamine in 0.2 M phosphate buffer (pH 8.0), nonspecific enzyme was removed from column by washing alternately in 0.2 M phosphate buffer (pH 8.0) containing 1 M NaCl, and in 0.2 M acetate buffer (pH 4.0) containing 1 M NaCl.

Determination of trypsin antibodies. The amount of antibodies in rabbit serum immunized with a suspension of tissue antigens of pancreas was determined in the column (0.5×10 cm) packed with Spheron P-1000 with immobilized trypsin. To assay the quantity of antibodies, 1 mg of serum was introduced into the column with an immunosorbent. After 15 min of incubation with a physiological solution at pH 7.4, the nonspecific proteins were washed out, and at pH 3.2 the specific ones. The quantity of protein was determined at the absorption of 280 nm.

Results and discussion

Products found as a result of the activation reaction of HMG with TCT and DCT are shown in Scheme.

The presence of two or three reactive chlorine atoms in the molecule of chlorine-substituted triazines leads to the formation of monosubstituted ester of triazine or the substitution for the second chlorine atom in the molecule of triazine, thus giving rise to the formation of absolutely undesirable intermolecular bonds according to scheme II.

Data in the Table show that reaction of HMG with TCT proceeds mainly according to scheme II, i. e., the second chlorine atom is sufficiently reactive and the reaction products obtained contain additional intermolecular bonds. In the reaction of HMG with DCT the substitution reaction proceeds mainly according to scheme I, i. e., the second chlorine atom is considerably less reactive and the formation of additional intermolecular bonds is insignificant. Varying reaction conditions, i. e., decreasing the reaction time, it is possible to obtain a gel without additional bonds.

As can be seen from Table, trypsin bound to Spheron C-1000 is of low reactivity and therefore will not find use in practice.

In order to optimize the binding conditions of biologically active molecules to HMG activated with chlorine derivates of triazine, trypsin has been used as a model compound.

Reaction of chlorine derivates of triazine with HMG

N	Triazine	N, %	Cl, %	Activity of trypsin, U/g
1	TCT	5.15	4.66	85
2	TCT	3.01	2.80	90
3	TCT	3.66	4.20	69.60
4	DCT	2.21	1.20	84.00
5	DCT	2.30	1.20	160.00
6	DCT	0.56	0.39	100.00
7	DCT	0.20	0.15	70.00
8	DCT	8.47	3.05	81.20
9	DCT	9.90	4.38	73.90
10	DCT	7.10	2.34	91.30
11*	—	—	—	42.00

* Spheron C-1000 for binding trypsin containing *p*-nitrophenyl-ester groups.




Fig. 1. Dependence of esterase activity of immobilized trypsin on the binding time: 1 — Sp-CN (Spheron P-1000, activated with TCT), 2 — Sp-ACN Spheron P-1000, activated with DCT).




Fig. 2. Dependence of esterase activity of Spheron-bonded trypsin on the amount of the enzyme and on the degree of HMG substitution (according to the chlorine content): 1—3 — Sp-ACN, 4 — Sp-CN.




Fig. 3. Dependence of esterase activity of immobilized trypsin on pH of the medium: 1 — Sp-CN, 2 — Sp-ACN.




Fig. 4. Dependence of esterase activity of immobilized trypsin on the chlorine content of Spheron.

vely less substituted HMG containing about 1% chlorine. The activity of bound trypsin decreases considerably with the increase of the chlorine content in HMG from 1 to 5%. The amount of the enzyme bound to the polymer matrix increases with the increase in the reaction time and the degree of the HMG substitution. We have demonstrated that the comparative activity of bound trypsin depends on the amount of trypsin bound to the polymer matrix so that the relative activity of the bound enzyme decreases considerably with an increase in its amount (Fig. 6).

All of the samples of HMG which have been activated with either TCT or DCT have excellent esterase activity that they retain during several




Fig. 5. Dependence of the relative esterase activity of Spheron P-1000 with immobilized trypsin on the amount of the bound trypsin.





Fig. 6. Esterase activity, pH optimum of Spheron with trypsin.

Fig. 7. The elution chromatogramme for antibodies in column with trypsin.

months and that does not disappear after a 20-cycle work.

At the same time, the proteolytic activity of trypsin bound to HMG with the aid of TCT approaches zero. It serves as an evidence of the additional bonds by activation of HMG, due to which the pores of HMG become narrower. These pores, although accessible to the comparatively small molecule of trypsin and its low substrate BAEE, are no more accessible to the large molecules of casein. The proteolytic activity of samples of trypsin bound to HMG with the aid of DCT is within 100 and 160 μm NaOH per 5 min (i. e., 5 to 9% relatively) and is stable enough.



Our data indicate the capacity of Spheron P-1000 activated with DCT to be lower than that of other carriers used for the quantitative assay of antibodies in rabbit serum. But taking into consideration the high properties of Spheron P-1000 under column conditions, its low capacity does not prevent using Spheron P-1000 in affinity chromatography of antibodies. The curve for elution in Fig. 7 shows the possibility of using immobilized trypsin to isolate antibodies of trypsin from rabbit serum immunized with a suspension of tissue antigens of pancreas.

Our investigations led to the conclusion that the binding of biological molecules to HMG with the aid of DCT may be considered possible, with the necessity of using preparations of low-degree substitution.

Summary

1. The activation reaction of hydroxyalkyl methacrylate gel Spheron P-1000 with chlorine derivatives of triazine, i. e., TCT and DCT, has been investigated.
2. Optimal conditions of binding trypsin to HMG-activated chlorine derivatives of triazine have been found.
3. Samples of the bound trypsin with a stable activity have been obtained by activating HMG with TCT and DCT.
4. The use of immobilized trypsin in the isolation of trypsin antibody from rabbit serum immunized with a suspension of tissue antigens of pancreas has been studied.

REFERENCES

1. Turková, O., Hubalková, O., Kriváková, M., Coupek, L. Affinity chromatography on hydroxyalkyl methacrylate gels. — *Biochim. Biophys. Acta*, 1973, v. 322, p. 1—9.
2. Fedoseyev, V., Pank, M., Murel, A., Heinlo, H., Kirret, O. Optimal bounding conditions and properties of trypsin samples immobilized on Sephadex by 2-amino-4,6-dichloro-S-triazine. — *ENSV TA Toim. Keem. Geol.*, 1977, kd. 26, nr. 4, lk. 326—329.
3. Kay, G., Lilly, M. D. The chemical attachment of chymotrypsin to water-insoluble polymers using 2-amino-4,6-dichloro-S-triazine. — *Biochim. Biophys. Acta*, 1970, v. 198, p. 276—285.

*Academy of Sciences of the Estonian SSR,
Institute of Chemistry*

Received
November 28, 1978

*Tiina LINNAS, Aili KOGERMAN,
H. HEINLO, O. KIRRET*

HUDROKSÜÜLALKÜÜLMETAKRÜLAATGEELIDE REAKTSIOON TRIASIINI KLOORDERIVAATIDEGA

Artiklis on käsitletud trüpsiini immobiliseerimist *Spheron* P-1000-le triasiini klooriderivaatide (2,4,6-trikloortriasiini ja 2-amino-4,6-dikloortriasiini) abil ning immobiliseeritud trüpsiini kasutamist trüpsiini antikehade eraldamiseks küülikute verest, keda eelnevalt on immuniseeritud pankrease ekstrakti vastaste antigeenidega.

*Тийна ЛИННАС, Айли КОГЕРМАН,
Х. ХЕЙНЛО, О. КИРРЕТ*

РЕАКЦИЯ ГИДРОКСИАЛКИЛМЕТАКРИЛАТНЫХ ГЕЛЕЙ С ХЛОПРОИЗВОДНЫМИ ТРИАЗИНА

Установлено, что перспективным можно считать связывание биологически активных молекул с -ОН-группами марки *Spheron* P-1000 при применении менее ядовитых реагентов — 2-амино-4,6-дихлортриазина и 2,4,6-трихлортриазина. Показана возможность использования иммобилизованного трипсина для выделения антител трипсина из сыворотки крови кроликов, иммунизированных суспензией тканевых антигенов поджелудочной железы.