

УДК 574.64; 582.26/27 (261.24)

O. ROOTS, H. KUKK

POLYCHLORINATED BIPHENYLS AND CHLORORGANIC PESTICIDES IN ALGAE FROM THE BALTIC SEA

O. ROOTS, H. KUKK. KLOORORGAANILISTE PESTITSIIDIDE JA POLUKLOREERITUD BIFENÜÜLIDE SISALDUS LÄÄNEMERE VETIKATES

O. РООТС, Х. КУКК. СОДЕРЖАНИЕ ХЛОРОРГАНИЧЕСКИХ ПЕСТИЦИДОВ И ПОЛИХЛОРИРОВАННЫХ БИФЕНИЛОВ В ВОДОРОСЛЯХ БАЛТИЙСКОГО МОРЯ

(Presented by O. Eisen)

Polychlorinated biphenyls (PCB) and chlororganic pesticides attract attention, first of all, due to their long-term existence in the surrounding environment and their ability to accumulate in living organisms. In our previous papers we have analyzed PCB and DDT concentrations in the plankton [1, 2], fish [3, 4] and molluscs [5] of the Baltic Sea.

Macroalgae may be considered important indicators when studying the distribution, transformation and decomposition of chlororganic compounds in the ecosystem of the Baltic Sea. Since algae are not included in the monitoring programme of the Baltic Sea, their biphenyl content has been insufficiently studied so far [6].

Algae samples were collected on September 16—19, 1984, during the cruise of the research vessel "Aju-Dag". To eliminate the possible contamination of algae samples with ship's paint, samples were collected from a boat at the distance of one kilometre from the ship [7]. The technique applied for the determination of the chlororganic compounds is presented in [3]. Clophen A-50 served as a PCB standard.

To get a more detailed picture about the transition of substances in the sea the coefficients characterizing their bioaccumulation from the water into algae were calculated (the Table). In the water of the open Baltic (Ariste Bay) and the Gulf of Finland (Kolga Bay) the concentrations of PCB and the summary DDT were calculated as 6.9 and 0.36 ng/l, and 4.3 and 0.13 ng/l, respectively (in the air above the open part of the sea PCB concentrations did not exceed 0.13 ng/m³, and those of the summary DDT 0.09 ng/m³). In most cases (with the exception of DDT in Kolga Bay) the coefficients obtained proved to be lower than the earlier ones in fish and plankton [8]. The concentrations of chlororganic pesticides determined in the water calls for careful application because in the study area they may show considerable diurnal variations depending on the water temperature, salinity, the content of plankton in the water, etc. [4]. The concentration of the summary DDT, as a rule, does not exceed 0.1 ng/l. In shallow sea areas one may observe the so-called secondary pollution where under certain conditions PCB may accumulate from bottom sediments into water and even re-enter the atmosphere [9].

Concentration of chlororganic pesticides and polychlorinated biphenyls in algae from the Baltic Sea, June 1984 (on wet weight (I) and on lipid weight (II) basis, $\mu\text{g}/\text{kg}$)

Biological object	Station	Lindan	p,p'/DDE	p,p'/DDD	p,p'/DDT	Σ DDT	PCB	Coef. bio-accum.		
								DDT	PCB	
<i>Fucus vesiculosus</i>	Ariste Bay	I	0.06	1.3	—	1.7	3.2	12.9	9000	1870
		II	4.3	93	—	124	226	920		
<i>Furcellaria lumbricalis</i>	"	I	0.05	0.6	—	2.3	3.0	9.5	8300	1380
		II	2.8	34	—	129	167	528		
<i>Ceramium rubrum</i>	"	I	0.12	0.3	—	—	0.3	8.7	830	1260
		II	40.0	103	0.6	—	113	2900		
<i>Fucus vesiculosus</i>	Kolga Bay	I	0.08	1.0	106	4.2	6.0	9.4	46150	2170
		II	13.3	158	—	700	992	1567		
<i>Cladophora glomerata</i>	"	I	0.23	3.3	—	5.5	9.2	37.9	70770	8750
		II	16.4	236	—	395	654	2707		
<i>Ectocarpus confervoides</i>	"	I	0.46	1.0	—	6.2	7.3	12.0	56150	2770
		II	65.7	141	—	886	1041	1714		
<i>Pilayella littoralis</i>	"	I	0.05	1.5	—	—	1.6	7.3	12310	1690
		II	8.3	248	—	—	273	1216		
<i>Dictyosiphon foeniculaceus</i> + <i>Ceramium tenuicorne</i>	"	I	0.006	—	—	—	—	—	—	—
		II	4.6	74	—	—	81	677	8460	2030

The above-mentioned shows that the results obtained are difficult to compare since the lipid concentrations may vary with different species of algae as well. The lowest PCB concentrations (on lipids) were determined in *Furcellaria lumbricalis* (an algae species of commercial importance in Estonia) and in *Dictyosiphon foeniculaceus* + *Ceramium tenuicorne*. In the Baltic Sea *Furcellaria* has revealed the lowest benzo(a)pyrene content as compared with other algae species [10]. PCB and DDT concentrations vary also in the samples of algae of the same species collected from different areas. PCB concentrations determined in *Fucus vesiculosus* proved to be lower than those determined by M. Olsson et al. in 1969 [6].

REFERENCES

1. Роотс О. О., Пейкре Э. А. О содержании устойчивых бифенилов и пестицидов в зоопланктоне Балтийского моря. — Водные ресурсы, 1981, № 6, 182—186.
2. Roots, O., Peikre, E. The study of the Baltic Sea zooplankton pollution with PCB and chlorinated pesticides during the 10th cruise of the R/V «Aju-Dag». — In: The Investigation and Modelling of Processes in the Baltic Sea (ed. A. Aitsam). Tallinn, 1981, Part II, 131—137.
3. Роотс О., Пейкре Э. О содержании полихлорированных бифенилов и хлорорганических пестицидов в рыбах Балтийского моря. — Изв. АН ЭССР. Хим., 1978, 27, № 3, 193—196.
4. Roots, O., Peikre, E. PCB and chlororganic pesticides in the Baltic Sea environment. — In: Proc. XIV Conf. of the Baltic Oceanographers. Gdynia, 1984, 2, 718—732.

5. *Roots, O.* PCB and chlororganic pesticides in molluscs of the Tallinn Bay. — In: The Investigation and Modelling of Processes in the Baltic Sea (ed. A. Aitsam). Tallinn, 1981, Part II, 137—141.
6. *Olsson, M., Jensen, S., Renberg, L.* PCB in coastal areas of the Baltic. — In: PCB II Conference. Stockholm, 1973, 59—68.
7. *Jensen, S., Renberg, L., Olsson, M.* PCB contamination from boat bottom paint and levels of PCB in plankton outside a polluted area. — Nature (London), 1972, N 240, 358—360.
8. *Roots, O., Peikre, E.* PCB and chlorinated pesticides in the ecosystem of the Baltic Sea. — In: Proc. XIII Conf. of the Baltic Oceanographers. Helsinki, 1982, 2, 604—616.
9. *Södergren, A., Larsson, P.* Transport of PCBs in aquatic laboratory model ecosystems from sediments to the atmosphere via the surface microlayer. — Ambio, 1982, N 11, 41—45.
10. *Veldre, I., Itra, A., Kukk, H., Paalme, L.* Benzo(a)pireeni sisaldusest vetikates. — ENSV TA Toim. Biol., 1981, 30, N 1, 51—53.

*Institute of Applied Geophysics,
Baltic Branch*

Received
Feb. 5, 1988

*Tallinn Department
of the Baltic Fishery Research Institute*

Polychlorinated biphenyls (PCB) and chlororganic pesticides in molluscs of the Tallinn Bay. — In: The Investigation and Modelling of Processes in the Baltic Sea (ed. A. Aitsam). Tallinn, 1981, Part II, 137—141.

Macrosalgae may be considered important indicators when studying the distribution, transformation and biodegradation of pollutants in aquatic environments. The results obtained in the present study show that the highest concentrations of PCB and DDT were found in the water of the open Baltic (Ariste Bay) and the Gulf of Finland (Sõiga Bay) and the concentrations of PCB and DDT were calculated as 6.6 and 0.6 ng/l, respectively. The highest concentrations of PCB and DDT were found in the water of the open Baltic (Ariste Bay) and the Gulf of Finland (Sõiga Bay) and the concentrations of PCB and DDT were calculated as 6.6 and 0.6 ng/l, respectively.

The investigation and modeling of PCB and chlororganic pesticides in molluscs of the Tallinn Bay. — In: The Investigation and Modelling of Processes in the Baltic Sea (ed. A. Aitsam). Tallinn, 1981, Part II, 137—141.

The investigation and modeling of PCB and chlororganic pesticides in molluscs of the Tallinn Bay. — In: The Investigation and Modelling of Processes in the Baltic Sea (ed. A. Aitsam). Tallinn, 1981, Part II, 137—141.