

## UPPER CAMBRIAN BIOSTRATIGRAPHY OF ESTONIA

Kaisa MENS,<sup>a</sup> Viive VIIRA,<sup>a</sup> Ivo PAALITS,<sup>b</sup> and Ivar PUURA<sup>a</sup>

<sup>a</sup> Eesti Teaduste Akadeemia Geoloogia Instituut (Institute of Geology, Estonian Academy of Sciences), Estonia pst. 7, EE-0100 Tallinn, Eesti (Estonia)

<sup>b</sup> Tartu Ülikooli Geoloogia Instituut (Institute of Geology, Tartu University), Vanemuise 46, EE-2400 Tartu, Eesti (Estonia)

Present address: Institut für Geologie und Paläontologie, Technische Universität Berlin (Technical University of Berlin, Institute of Geology and Paleontology), Ackerstraße 79—76, Sekr. ACK 14, Berlin 13355, Deutschland (Germany)

Presented by D. Kaljo

Received April 2, 1993; accepted May 28, 1993

**Abstract.** In Estonia the Upper Cambrian is represented by siliciclastic rocks, mainly quartzose sand- and siltstones containing thin interlayers of greenish-grey clays in the lower and dark kerogenous argillites in the upper part of the section. The Upper Cambrian of Estonia is subdivided into four formations: Petseri, Ülgase, Tsitre, and Kallavere.

The biostratigraphic subdivision is based on three fossil groups—acritarchs, conodonts, and lingulates. Five acritarch-, four conodont-, and at least three lingulate-based biostratigraphic units are distinguished, which may be used in the Upper Cambrian stratigraphic scale of Estonia. Subdivision of the lower part of the section is more reliable on acritarchs and that of the upper part, on conodonts.

Using the succession of acritarchs and conodonts the Petseri and Ülgase formations are approximately correlated with the *Olenus* Zone and the latter also with the *Parabolina spinulosa* Zone (lower part). The Tsitre Formation (lower part) corresponds to the *P. spinulosa* (upper part) and *Leptoplastus* zones, its uppermost part to the *Peltura scarabaeoides* Zone and the lower part of the Kallavere Formation can be referred to *Acerocare* Zone.

**Key words:** acritarchs, conodonts, lingulates, Upper Cambrian, Estonia.

### INTRODUCTION

In the framework of biostratigraphic studies of the Cambrian—Ordovician boundary beds in Estonia and neighbouring areas, a number of key sections have been described in detail during the past 10 years with the documentation of the distribution of acritarchs, conodonts, and lingulate brachiopods. Conodont zonation has also contributed to detailed biostratigraphic correlation of the Upper Cambrian deposits exposed in the outcrops along the Baltic—Ladoga clint (Kaljo et al., 1986; Попов et al., 1989). Alongside conodonts, lingulate brachiopods have proved to be useful for basinwide correlations, particularly between Estonian and Swedish sections (Puura & Holmer, 1993). Acritarchs, which occur only in thin argillaceous interbeds of the sandstone sequence, serve as a basis for a detailed biostratigraphic subdivision of sections and for correlation with distant sequences worldwide (Волкова, 1990; Paalits, 1992a, 1992b).

Only part of the results of this research has been published, mostly as detailed descriptions of separate key sections (e.g. Хейнсалю et al., 1987, 1991; Kaljo et al., 1988; Mens et al., 1989).

The purpose of the present paper is to document the co-occurrence of acritarchs, conodonts, and lingulates in Estonian Upper Cambrian sections and to discuss the relationships of their biounits. A tentative correlation of acritarch, conodont, and lingulate biounits with Scandinavian trilobite zones is presented.

## GEOLOGICAL SETTING

In the present paper the Upper Cambrian is considered in its traditional extent for the East European Platform, i.e. from the base of the *Agnostus pisiformis* Zone to the top of the *Acerocare* Zone. As trilobites are lacking in Estonia and *Cordylodus proavus* is documented in the Nærnes section, Norway, together with trilobites of the *Acerocare* Zone (Bruton et al., 1988), we have included also the *C. proavus* Zone level into our discussion.




Fig. 1. The distribution, thickness, and facies of the Upper Cambrian in the north-western part of the East European Platform.

1 — thickness of the corresponding deposits (up to the base of the *Cordylodus* succession); the question mark shows an undetermined Cambrian or Ordovician age of deposits; 2 — recent erosional boundary of the Upper Cambrian; 3 — isopachytes; 4 — boundary of a facies belt.

The Upper Cambrian rocks, which have been preserved from post-Cambrian erosion, are widely distributed in the north-western part of the East European Platform ranging as a sublatitudinal zone from the Oslo graben to the Moscow syncline (Fig. 1). They are exposed along the Baltic—Ladoga clint area and in the river valleys intersecting it, occurring in the form of isolated patches also in the Oslo graben, South and Central Sweden, and are often allochthonously lying near the north-western margin of the Platform. In the rest of the territory, the Upper Cambrian rocks are lying at a depth from a few metres (e.g. northern Estonia and southern Scandinavia) to two thousand metres (central part of the Moscow syncline).

From the west to the east, the following lateral lithofacies sequences can be observed. In the west, approximately up to the Prabuty—Öland—Närke line, alum shales are prevailing. Eastwards, there occurs a facies

of sand- and siltstones, with rare thin interlayers of so-called *Dictyonema* shale, clay and sandstones with lenses of lingulate brachiopod valves and debris. East of the Ladoga—Porhov line, the Upper Cambrian is represented by siltstones and clays, more rarely by sandstones, while organic-rich rocks are absent (Fig. 1).

The sections of these three areas, tentatively distinguished as western, transitional, and eastern facies belts, differ not only in lithology and thickness, but also in stratigraphic completeness (see Mens et al., 1990) and paleontological evidences.

The Upper Cambrian biostratigraphy of the western facies belt is mainly based on trilobites and their zonation serves as a standard for platform correlation and a basis for interregional correlation (Westergård, 1922). From these sections of the transitional belts no trilobites have been found whereas the sections of the eastern facies belt have yielded only a few specimens. In the transitional belts the Upper Cambrian biostratigraphy is based on the other groups occurring in abundance, namely, acritarchs, conodonts, and lingulate brachiopods.

The Upper Cambrian sequence of Estonia is exposed in numerous outcrops and boring cores (Fig. 2). This has allowed of detailed investigation of the corresponding deposits.




Fig. 2. Location of the main outcrops (triangles) and borings (solid circles) mentioned in the text.

In Estonia the strata considered as the Upper Cambrian are siliciclastic, dominated by silt- and sandstones less than 20 m in thickness (Fig. 1). Argillaceous rocks are limited, forming greenish-grey clayey interlayers within light-coloured coarse-grained deposits while in the uppermost part of the section they are often represented by their kerogen-bearing variety, the *Dictyonema* shales.

The Upper Cambrian succession is very condensed and interrupted by several minor as well as major hiatuses. The presence of gaps is supported by petrographic evidences and by the absence of one or some biounits in the section.

The Upper Cambrian lies mainly on Lower Cambrian silt- and sandstones and only locally on Middle Cambrian rocks. It is overlain by Tremadocian siliciclastics, in places by younger Ordovician limestones.

According to the stratigraphic subdivision, the Upper Cambrian

includes the Petseri, Ülgase, and Tsitre formations and a part of the Kallavere Formation. The Petseri Formation, known only in the subsurface, is distributed in south-eastern Estonia (see Волкова et al., 1981). The other three formations occur more widely in northern Estonia (Kaljo et al., 1986; Mens et al., 1990; etc.).

**Acritarchs** have been studied from the Tõnismägi, Mäekalda, Suhkrumägi, Ülgase, Valkla, Turjekelder, Vihula, and Saka outcrops and from the boring cores M-9, M-36, M-46, M-56, M-72, M-77, P-2162, Hino, Mehikoorma, Petseri, and Põlva (Fig. 2).

The taxonomic diversity and stratigraphic as well as relatively wide lateral distribution of acritarchs have enabled to establish the guide assemblages within the Upper Cambrian succession of the East European Platform including six acritarch assemblages (BK 1, BK 2, BK 3, BK 4A, BK 4B, BK 5) by Volkova (Волкова, 1990). Three similar assemblages (BK 3, BK 4B, BK 5) have been established in the Estonian Upper Cambrian while the assemblage BK 2 can be divided into two independent parts with key species *?Dasydiacrodium setuensis* and *Leiofusa stoumonensis* for the lower and *Impluviculus multiangularis*—*?Veryhachium dumontii* for the upper one. The assemblages BK 1 and BK 4A, corresponding respectively to the *A. pisiformis* Zone and to the lower part of the *Peltura* Superzone, have not been determined in Estonia. Thus, acritarchs provide a valuable tool for the biostratigraphic subdivision of the Estonian Upper Cambrian and for correlating it with that of the eastern and western facies belts. As acritarchs are more common and their assemblages more diverse in clayey and argillaceous rocks occurring as thin interlayers between sand- and siltstones, it is impossible to draw exact boundaries between acritarch biounits.

The oldest Upper Cambrian assemblage recorded from the rocks of the Petseri Formation and represented by *Cristallinium cambriense* (Slavikova), *Cymatiogalea dentalea* Paalits, *C. virgulta* Martin, *?Dasydiacrodium setuensis* Paalits, *Leiofusa stoumonensis* Vanguetstaine, *Stelliferidium* aff. *certinulum* (Deunff) Deunff, Gorka & Rauscher, *Timojeevia lancaerae* (Cramer & Diez) Vanguetstaine, *T. phosphoritica* Vanguetstaine, *Veryhachium incus* Paalits, *Vulcanisphaera turbata* Martin, and species of the genera *Poikilofusa*, *Timojeevia*, and *Micrhystridium* (Волкова, 1990; Paalits, 1992b), has not been found anywhere else in Estonia.

Based on the occurrence of the genera *Stelliferidium*, *Cymatiogalea*, *Leiofusa*, and *Veryhachium*, which appeared in the *Olenus* time (Potter, 1974; Downie, 1984), and the absence of the *Impluviculus* species, the deposits yielding this acritarch assemblage are considered as a stratigraphic equivalent of the lower and/or middle parts of the *Olenus* Zone.

The next acritarch assemblage is similar to the one described, but differs in the presence of the representatives of the genus *Impluviculus* and in the lack or restricted distribution of typical Middle Cambrian species (Волкова, 1982). In all the studied sections this assemblage occurs in the Ülgase Formation. As the appearance of the genus *Impluviculus* has been correlated with the uppermost part of the *Olenus* Zone (Downie, 1984) and with the lowermost part of the *P. spinulosa* Zone (Martin & Dean, 1988), we regard the Ülgase Formation tentatively as a stratigraphic equivalent of the uppermost *Olenus* and the lowermost *Parabolina* zones (Table).

The third Upper Cambrian assemblage, containing *Trunculumarinum revinium* (Vang.) Loeblich et Tappan, *Dasydiacrodium caudatum* Vanguetstaine, *D. obsenum* Martin, *Leiofusa stoumonensis* Vanguetstaine, *Veryhachium dumontii* Vanguetstaine, *Cymatiogalea wironia* Paalits, *C.*

Suggested correlation of the Upper Cambrian biostratigraphic and lithostratigraphic units of Estonia and their relationship with trilobite zones

Trilobite zones	Conodont zones & subzones	Lingulate zones	Key species of acritarchs	Lithostratigraphic units	
				N and NW Estonia	SE Estonia
<i>Boeckaspis hirsuta</i>	<i>C. lindstromi</i> <i>C. intermedius</i>	<i>O. apollinis</i>	<i>A. echinatum</i> <i>A. striatum</i>	Kallavere Fm. (part)	Kallavere Fm. (part)
<i>Acerocare</i>	<i>C. proavus</i> <i>C. andresi</i>	<i>U. ingrca</i>	<i>A. angustum</i> <i>D. palmatilobum</i> <i>I. angulata</i> <i>O. rossicum</i> <i>E. armillata</i> <i>V. cervinacornua</i>		
<i>Peltura scarabaeoides</i>	<i>Proconodontus</i>				
<i>Peltura minor</i>			?	Tsitre Fm.	
<i>Protopeltura praecursor</i>	?	<i>U. convexa</i>			
<i>Leptoplastus</i>			<i>T. revinium</i> <i>D. caudatum</i>		
<i>Parabolina spinulosa</i>					
<i>Olenus</i>	<i>W. bicuspidata</i>	<i>U. inornata</i>	<i>I. multiangularis</i> ? <i>V. dumontii</i>	Ülgase Fm.	
<i>Agnostus pisiformis</i>			? <i>D. setuensis</i> <i>L. stauntonensis</i>		Petseri Fm.

O

-?

€

Westergaardodina

*dentalea* Paalits, *C. aff. virgulta* Martin, *C. sp. sp.*, *Goniosphaeridium aff. tuberatum* (Downie) Wolf, *G. aff. dentatum* (Timofeev) Rauscher, *G. sp. sp.*, *Stelliferidium cortinulum* (Deunff) Deunff, Gorka & Rauscher, *S. aff. pseudoornatum* Pittau, *Timofeevia phosphoritica* Vanguestaine, *T. estonica* Volkova, *Vulcanisphaera turbata* Martin, *V. sp.*, *Actinodissus sp.*, *Cristallinium sp.*, corresponds to the microflora A<sub>4</sub>, i.e. *T. revinium*—*D. caudatum* assemblage *sensu* Martin & Dean (1988) and to assemblage BK-3 *sensu* Volkova (Волкова, 1990). In northern Estonia, this assemblage occurs in the lowermost part of the Tsitre Formation found only in boring cores M-9, depth 134.5—140.6 m; M-46, depth 38.7—40.5 m; M-72, depth 112.8—119.6 m (Paalits, 1992a.; Волкова, 1990). The tentative correlation of this assemblage with the trilobite zones is based on *Trunculumarinum revinium* and *Dasydiacrodium caudatum* known as the species with a limited stratigraphic range from the uppermost *P. spinulosa* Zone through the whole *Leptoplastus* Zone (Table).

The next acritarch assemblage, showing high taxonomic diversity and variation in different sections, contains a significant amount of diacroids and maybe endemic forms. On the East European Platform this assemblage was first described from the upper part of the Ladoga Formation, St. Petersburg area (Волкова & Голуб, 1985), and is referred to as BK-4Б (Волкова, 1990). It is characterized by the prevalence of *Acanthodiacrodium timofeevii* Volkova et Golub, *Ladogiella rotundiformis* Volkova et Golub, *Calyziella izhoriensis* Volkova et Golub, *Izhoria angulata* Volkova et Golub, *Nellia longiscula* Volkova et Golub, *Schizodiacrodium fibrosum* Volkova et Golub, *Dasydiacrodium palmatilobum* Timofeev, *Elenia armillata* (Vand.) Volkova, in some samples by *Ooidium timofeevii* Loeblich and *O. rossicum* Timofeev. In Estonia this acritarch assemblage is found from the upper part of the Tsitre Formation of the Valkla, Turjekelder, and Saka outcrops and from the boring cores P-2162, depth 115.4—115.6 m, and M-56, depth 138.6—147.4 m, together with the conodonts of the *Proconodontus* Subzone. These acritarchs have also been found in the *Cordylodus andresi* Zone of the Vihula section (Волкова, 1990). A relatively similar assemblage together with the trilobites of the *Peltura scarabaeoides* Zone has been determined from the Degerhamn section of southern Öland (di Milia et al., 1989).

The youngest Upper Cambrian acritarch assemblage which, as the above one, contains a large number of diacroids (Волкова, 1989, 1990), can be distinguished by the appearance of *Acanthodiacrodium angustum* (Downie) Combaz and *Dicrodiacrodium ramusculosum* (Combaz) Volkova. It is found together with conodonts of the *C. proavus* Zone from the lower part of the Kallavere Formation of the Tõnismägi, Mäekalda, Suhkrumägi, and Vihula outcrops and from the boring core M-9, depth 131.1 m.

**Conodonts** have been studied from a number of outcrops and boring cores embracing the Cambrian—Ordovician boundary beds. Late Cambrian conodonts have been found in the Mäekalda, Suhkrumägi, Ülgase, Valkla, Turjekelder, Vihula, Toolse, and Saka outcrops and from the Kidaste and M-9 boreholes (Fig. 2). The number of specimens is small, representing mostly the genera *Phakelodus*, *Furnishina*, *Prooneotodus*, and *Westergaardodina*. Eoconodonts (*Proconodontus*, *Eoconodontus*, and *Cordylodus*) have been found only in the uppermost Upper Cambrian.

Here we accept the conodont zonation proposed by Viira and Sergeeva for the correlation of the Cambrian—Ordovician boundary beds along the Baltic—Ladoga clint area (see Боровко & Сергеева, 1985; Kaljo et al., 1986). According to this zonation, the Upper Cambrian sequence is subdivided as follows, starting from below: the

## ÜLGASE

Acritarchs	Conodonts	Lingulates
<i>A. angustum</i> (1.7–2.05 m)	<i>C. proavus</i> (1.3–2.05 m)	<i>U. ingrlica</i> <i>S. celatus</i> , <i>K. buchi</i> (1.7–4.8 m)
	<i>Prooneotodus?</i> sp. (2.3–3.2 m)	
	<i>P. perforata</i> , <i>W. cf. bicus-</i> <i>pidata</i> (3.2–4.1 m)	
<i>I. multiangu-</i> <i>laris</i> (4.8–5.1 m)	<i>P. tenuis</i> <i>Furnishina</i> sp. (4.8–5.1 m)	<i>U. inornata</i> (4.8–5.1 m)

Depth data from Хейнсаалу et al., 1987

## M-9

Acritarchs	Conodonts	Lingulates
<i>A. angustum</i> (131 m)	<i>C. proavus</i> (131 m)	
<i>T. revinium</i> (134.5–146 m)		

Depth data from Волкова, 1989

## TURJEKELDER

Acritarchs	Conodonts	Lingulates
<i>D. palmati-</i> <i>lobum</i> , <i>E. armillata</i> , <i>O. rossicum</i> (4.8–8.4 m)	<i>C. andresi</i> (7.4–8.4 m)	<i>U. ingrlica</i> , <i>S. celatus</i> , <i>K. buchi</i> (4.7–7.0 m)
	<i>P. cf. gallaini</i> <i>W. bicuspidata</i> (4.7–6.0 m)	
<i>I. multiangu-</i> <i>laris</i> (0.5–4.7 m)	<i>P. tenuis</i> , <i>F. furnishi</i> (0.0–4.7 m)	<i>U. inornata</i> (0.0–4.7 m)

Depth data from Kaljo et al., 1986

## M-72

Acritarchs	Conodonts	Lingulates
<i>T. revinium</i> (112.8–116 m)		<i>U. convexa</i> <i>S. celatus</i> (115–116 m)

Depth data from Paalits, 1992a

## VIHULA

Acritarchs	Conodonts	Lingulates
<i>A. angustum</i> (1.7 m)	<i>C. proavus</i> (1.5–5.1 m)	<i>H. ladogensis</i> (2.3 m)
<i>D. palmati-</i> <i>lobum</i> <i>I. angulata</i> (0.1–1.5 m)	<i>C. andresi</i> (0.1–1.5 m)	<i>U. ingrlica</i> , <i>S. celatus</i> , <i>K. buchi</i> (0.0–0.4 m)

Depth data from Попов et al., 1989

## P-2162

Acritarchs	Conodonts	Lingulates
<i>V. cervina-</i> <i>cornia</i> (115.4–115.6 m)		<i>O. apollinis</i> (113 m)
		<i>U. convexa</i> (114–115.3 m)

*Westergaardodina* Zone with *W. bicuspidata*, *W. moessenbergensis*, and *Proconodontus* subzones, and the *Cordylodus andresi* and *C. proavus* zones.

The *Westergaardodina bicuspidata* Subzone is distinguished by the species *Phakelodus tenuis* (Müller), *Furnishina furnishi* Müller, *F. alata* Szaniawski, and *Prooneotodus terashimai* (Nogami). The zonal species *W. bicuspidata* is very rare. The conodont assemblage of the *W. bicuspidata* Subzone has been found from the Ülgase Formation of the Mäekalda, Suhkrumägi, Ülgase, Jägala, Valkla, and Turjekelder outcrops and boring core M-9 (Боровко & Сергеева, 1985; Хейнсалю et al., 1987; Kaljo et al., 1986; Mens et al., 1989). As provisionally correlated with the trilobite zonation, the *W. bicuspidata* Zone would correspond to the *Agnostus pisiformis*, *Olenus*, and the lowermost *Parabolina spinulosa* zones (Боровко & Сергеева, 1985).

The *Westergaardodina moessenbergensis* Subzone has not yet been established in Estonia (Kaljo et al., 1986).

The *Proconodontus* Subzone is distinguished by the occurrence of *Proconodontus primitivus* (Müller), *Prooneotodus* cf. *gallatini* (Müller), and *Problematoconites perforata* Müller appearing in the eastern sections of the Leningrad Region already in the *Westergaardodina moessenbergensis* Subzone. This conodont assemblage has been found from the Tsitre and lowermost Kallavere formations of the Ülgase, Turjekelder, and Saka sections. The thickness of the rocks corresponding to this zone is up to 2 m (Kaljo et al., 1986; Хейнсалю et al., 1987). Proceeding from the co-occurrence of the above-mentioned conodonts with such acritarchs as *Elenia armillata*, *Dacydiacrodium palmatilobum*, and *Ooidium rossicum* (Fig. 3), the *Proconodontus* Subzone is tentatively correlated with the *Peltura scarabaeoides* Zone.

The *Cordylodus andresi* Zone represents the beginning of the *Cordylodus* succession in many sections along the Baltic—Ladoga clint (Kaljo et al., 1986; Viira et al., 1987). The conodont assemblage of this zone is established in the lower, up to 2 m thick part of the Kallavere Formation of the Ülgase, Valkla, Turjekelder, Toolse, and Vihula outcrops and Kidaste borehole on Hiiumaa Island. Besides the zonal species, the assemblage of the *C. andresi* Zone contains *Eoconodontus notchpeakensis* (Miller), *Cordylodus viruanus* Viira et Sergeyeva, and the long-ranging species of an earlier appearance *Phakelodus tenuis*, *Furnishina furnishi*, *Westergaardodina bicuspidata*, *Prooneotodus* cf. *gallatini*, and *Muellerodus* sp. Outside the transitional facies belt, *C. andresi* has been found from the *Westergardia* Subzone of the *Acerocare* Zone on the island of Öland (Andres, 1981) and together with *Parabolina heres heres* from the Zhar-novez Region of northern Poland (Lendzion, pers. comm.).

The *Cordylodus proavus* Zone is established in numerous studied sections, mostly from the lower part of the Kallavere Formation. In addition to the index species *Eoconodontus notchpeakensis*, *Cordylodus andresi*, and long-ranging paraconodonts occur.

Recent studies of the Cambrian—Ordovician boundary interval in the Nærnes section, Norway, have shown that the first *Cordylodus proavus* appeared already on the level of the *Acerocare ecorne* Subzone, i.e. in the uppermost subzone of the *Acerocare* Zone (Bruton et al., 1988). Taking this into account, deposits of the *C. proavus* Zone and its sup-

Fig. 3. Co-occurrence of the key species of acritarchs, conodonts, and lingulates in selected Upper Cambrian sections of Estonia.

Indexes of stratigraphic units: O<sub>1</sub>—E<sub>3</sub>kl — Kallavere Formation, E<sub>3</sub>ts — Tsitre Formation, E<sub>3</sub>ül — Ülgase Formation. Double dashed line marks absence of one or some biounits.



posed stratigraphic equivalents are included within the Upper Cambrian (Table).

**Lingulate** brachiopods are represented in the Upper Cambrian of Estonia by lingulids and acrotretids. Their stratigraphic distribution has been studied in a number of sections including Tõnismägi, Mäekalda, Suhkrumägi, Ülgase, Valkla, Turjekelder, Vihula, and Saka outcrops and the Kidaste, Aiamaa, M-9, M-72, M-77, P-2162 boring cores, where acritarchs or conodonts have often been studied (Fig. 2).

Following the brachiopod zonation suggested by Popov and Khazanovitch (Попов et al., 1989) and adopted by Puura & Holmer (1993), the following four brachiopod zones can be distinguished, starting from below: the *Ungula inornata* Zone, the *Ungula convexa* Zone, the *Ungula ingrlica* Zone, and the *Obolus apollinis* Zone. The first three zones belong to the Upper Cambrian, while the last one can belong partly to the Ordovician, depending on the position of the lower boundary of the Ordovician System.

The *Ungula inornata* Zone yields, besides the zonal species, *Oepikites fragilis* Popov & Khazanovitch, *Ceratreta tanneri* (Metzger), and *Angulotreta postapicalis* Palmer. The lingulate assemblage of the *Ungula inornata* Zone has been found from the Ülgase Formation in the Tõnismägi, Mäekalda, Suhkrumägi, Iru, Ülgase, Valkla, and Turjekelder outcrops and M-9 and M-77 boring cores.

The *Ungula convexa* Zone yielding besides the zonal species *Oepikites triquetrus* Popov & Khazanovitch and *Keyserlingia reversa* (de Verneuil) was distinguished by Popov and Khazanovitch (Попов et al., 1989) only from the Leningrad Region and correlated with the lowermost Upper Ladoga Subformation. Recently, *Ungula convexa* has been found in the Tsitre Formation of Estonia from the Saka section and boring cores P-2162, depth 114—115.36 m; P-2163, depth 108.8—111.3 m; and M-72, depth 115—116 m, in the last case together with *Schmidtitites celatus*. Only *Oepikites triquetrus* has been determined from boring cores M-39, depth 165.2—168.7 m; M-77, depth 29.2 m.

The distributional data from boring core M-72 (Fig. 3) are interesting because *U. ingrlica* and *S. celatus* have been met together with an acritarch assemblage yielding *Trunculumarinium revinium* (Paalits, 1992a), which has a relatively narrow time span (Martin & Dean, 1988). On the ground of these data the lower boundary of the *U. convexa* Zone is lowered (Table) in comparison with the primary one (Попов et al., 1989).

The *Ungula ingrlica* Zone yielding *U. ingrlica* (Eichwald), *Schmidtitites celatus* (Volborth), *Keyserlingia buchii* (de Verneuil), and *Oepikites obtusus* (Mickwitz) corresponds to the uppermost part of the Tsitre Formation and the lowermost part of the Kallavere Formation of the Tõnismägi, Mäekalda, Suhkrumägi, Iru, Ülgase, Valkla, Turjekelder, Vihula, Äseri, and Saka outcrops and boring core M-77, depth 24.2—34.3 m. Its stratigraphic range is relatively clearly defined by conodont and acritarch co-occurrences.

The *Obolus apollinis* Zone with the lower boundary coinciding with that of the *Cordylodus proavus* conodont Zone in the Leningrad Region, yields the species *Obolus apollinis* (Eichwald) and *Helmerseniania ladogensis* (Jeremejew). These species are widely distributed in the Tosno Formation of the Leningrad Region, but rather rare in Estonia, being known from the boring cores of Kidaste, K-11, and K-19 on Hiiumaa Island and P-1555 and P-2162 of the Rakvere phosphate deposits area. *Helmerseniania ladogensis* has been found from K-11 core on Hiiumaa Island and reported from the Vihula section (Попов & Хазанович,

1989). Very often the finds of *O. apollinis* and/or *H. ladogensis* are related to the occurrences of *Cordylodus proavus* and *Acanthodiacrodium angustum*.

## DISCUSSION AND CONCLUSIONS

The studied three groups of fossils—acritarchs, conodonts, and lingulate brachiopods—complement one another in respect of biostratigraphic correlation. Therefore, the documentation of their co-occurrences in a series of Estonian Upper Cambrian sections (Fig. 3) gives useful information both for short- and long-distance biostratigraphic correlation.

The oldest Upper Cambrian lithostratigraphic unit of Estonia, the Petseri Formation, is at present characterized only by acritarchs including *Leiofusa stoumonensis* Vanguetaine and ?*Dasydiacrodium setuensis* Paalits (Table), which show early late Cambrian age.

The next lithostratigraphic unit, the Ulgase Formation, contains representatives of all the groups studied. Of the highest correlative value are *Impluviculus multiangularis*, *Westergardina bicuspidata*, and *Ungula inornata*, indicating the deposition of the corresponding sediments during the first half of the late Cambrian.

The lower part of the Tsitre Formation is characterized by an association of lingulates and acritarchs, the acritarch *Trunculumarinum revinium* having a short stratigraphic range. Conodonts have not been found as yet from this part of the sequence, which is revealed only in a few boring cores. The upper part of the Tsitre Formation, however, yields acritarchs, conodonts, and lingulates, including also conodonts of the *Proconodontus* Subzone (Fig. 3, Turjekelder Section).

The upper stratigraphic boundary of the acritarch assemblage containing *Dasydiacrodium palmatilobum*, *Ishoria angulata*, etc. coincides with that of the *Cordylodus andresi* Zone.

The most debatable is the position of the boundary between the *Ungula ingrlica* and *Obolus apollinis* zones in relation to the lower boundary of the *Cordylodus proavus* Zone. The acritarch assemblage with *Acanthodiacrodium angustum* appears at the base of the *Cordylodus proavus* Zone. In northern Estonian sequences *C. proavus* and *A. angustum* co-occur with *Ungula ingrlica*. *O. apollinis*, which is distributed in a sublatitudinal belt extending from the Leningrad Region to the Rakvere phosphate deposit area and Hiiumaa Island, is always known to appear above the base of the *C. proavus* Zone. Defining the boundary between *U. ingrlica* and *O. apollinis* zones by the first appearance of *O. apollinis*, we leave the question of its detailed correlation with the lower boundary of the *C. proavus* Zone open. Our provisional correlation provided in the Table shows the lower boundary of the *O. apollinis* Zone slightly below the top of the *Acerocare* Zone.

The tentative correlation of the Estonian Upper Cambrian biounits with Scandinavian trilobite zones is mostly based on acritarch correlation and only within the *Acerocare* Zone on conodonts. However, the direct correlation is possible only partly because of the lack of data on the distribution of acritarchs of the *Agnostus pisiformis*, *Leptoplastus*, *Protopeltura praecursor*, *Peltura minor*, and *Acerocare* zones in Scandinavia.

In all probability, stratigraphic analogues of the *Olenus*, *Parabolina spinulosa*, and *Peltura scarabaeoides* zones can be distinguished in Estonia. The *Cordylodus andresi* and *C. proavus* zones, respectively, correspond to the *Westergardia* and *A. ecorne* subzones of the *Acerocare* Zone.

The sandy lower part of the Petseri Formation has not yielded fossils. As the overlying clays in the middle part of this formation contain acritarchs of the *Olenus* Zone, the correlation of those barren sandstones with the *Agnostus pisiformis* Zone has been suggested (Mens et al., 1990).

The acritarch assemblage with *Trunculumarinum revinium* and *Dasydiacrodium caudatum* from the lower part of the Tsitre Formation, in addition to the upper part of the *P. spinulosa* Zone, is tentatively correlated with the *Leptoplastus* Zone.

## ACKNOWLEDGEMENTS

We thank Anne Noor for linguistic help. Technical support by Kaie Ronk and Urve Pohl is also gratefully acknowledged.

## REFERENCES

- Andres, D. 1981. Beziehungen zwischen kambrischen Conodonten und Eoconodonten. — Berliner Geowiss. Abh. A., 32, 19—31.
- Bruton, D. L., Koch, L., Repetski, J. E. 1988. The Nærnes section, Oslo Region, Norway: Trilobite, graptolite and conodont fossils reviewed. — Geol. Mag., 125, 4, 451—455.
- di Milia, A., Ribecai, C., Tongiorgi, M. 1989. Late Cambrian acritarchs from the *Peltura scarabaeoides* trilobite zone at Degerhamn (Öland, Sweden). — Palaeontographia Italica, LXXVI, 1—55.
- Downie, C. 1984. Acritarchs in British Stratigraphy. Geol. Soc. London. Special Report, 17, 261.
- Kaljo, D., Borovko, N., Heinsalu, H., Khazanovitch, K., Mens, K., Popov, L., Sergejeva, S., Sobolevskaya, R., Viira, V. 1986. The Cambrian—Ordovician boundary in the Baltic—Ladoga clint area (North Estonia and Leningrad Region, USSR). — Proc. Acad. Sci. ESSR. Geol., 35, 3, 97—108.
- Kaljo, D., Heinsalu, H., Mens, K., Puura, I., Viira, V. 1988. Cambrian—Ordovician boundary beds at Tõnismägi, Tallinn, North Estonia. — Geol. Mag., 125, 4, 457—463.
- Martin, F., Dean, W. T. 1988. Middle and Upper Cambrian acritarch and trilobite zonation at Manuels River and Random Island, eastern Newfoundland. — Bull. Geol. Surv. Canada, 381, 1—91.
- Mens, K., Bergström, J., Lenzion, K. 1990. The Cambrian System of the East European Platform (Correlation chart and explanatory notes). — Intern. Union of Geol. Sciences, 25, 78.
- Mens, K., Viira, V., Paalits, I., Puura, I. 1989. Cambrian—Ordovician boundary beds at Mäekalda (Tallinn, North Estonia). — Proc. Estonian Acad. Sci. Geol., 38, 3, 101—111.
- Paalits, I. 1992a. Upper Cambrian acritarchs from boring core M-72 of North Estonia. — Proc. Estonian Acad. Sci. Geol., 41, 1, 29—37.
- Paalits, I. 1992b. Upper Cambrian acritarchs from the Petseri Formation (East European Platform). — Tartu Ülikooli Toimetised, 956. Töid geoloogia alalt, XIII, 44—55.
- Potter, T. L. 1974. British Cambrian acritarchs—a preliminary account. — Rev. Palaeobot. Palynol. (Special issue on acritarchs), 18, 61, 62.
- Puura, I., Holmer, L. E. 1993. Lingulate brachiopods from the Cambrian—Ordovician boundary beds in Sweden.—Geologiska Föreningens i Stockholm Förhandlingar 115, pt. 3, 215—237.
- Viira, V., Sergejeva, S., Popov, L. 1987. Earliest representatives of the genus *Cordylodus* (Conodonta) from Cambro—Ordovician boundary beds of North Estonia and Leningrad Region. — Proc. Acad. Sci. ESSR. Geol., 36, 4, 145—153.

- Westergård, A. H. 1922. Sveriges Olenidskiffer. (The olenid shale of Sweden.) — Sver. Geol. Unders. Ca 18, 1—205.
- Боровко Н., Сергеева С. 1985. Конодонты верхнекембрийских отложений Балтийско-Ладожского глинта. — Изв. АН ЭССР. Геол., 34, 4, 125—129.
- Волкова Н. А. 1982. О возрасте юлгаской пачки на границе кембрия и ордовика в Эстонии. — Сов. геол., 9, 85—88.
- Волкова Н. А. 1989. Акритархи пограничных отложений кембрия и ордовика севера Эстонии. — Изв. АН СССР. Сер. геол., 7, 59—67.
- Волкова Н. А. 1990. Акритархи среднего и верхнего кембрия Восточно-Европейской платформы. Наука, Москва.
- Волкова Н. А., Голуб И. Н. 1985. Новые акритархи верхнего кембрия Ленинградской области (ладожская свита). — Палеонтол. ж., 4, 90—98.
- Волкова Н., Каяк К., Менс К., Пиррус Э. 1981. Новые данные о переходных слоях между кембрием и ордовиком на востоке Прибалтики. — Изв. АН ЭССР. Геол., 30, 2, 51—55.
- Попов Л. Е., Хазанович К. К. 1989. Лингулаты (беззамковые брахиоподы с фосфатнокальциевой раковиной). — Ип: Опорные разрезы и стратиграфия кембро-ордовикской фосфоритоносной оболовой толщи на северо-западе Русской платформы. Наука, Ленинград, 96—136.
- Попов Л. Е., Сергеева С. П., Хазанович К. К. 1989. Биостратиграфическое расчленение. — Ип: Опорные разрезы и стратиграфия кембро-ордовикской фосфоритоносной оболовой толщи на северо-западе Русской платформы. Наука, Ленинград, 85—95.
- Хейнсалу Х., Вийра В., Менс К., Оя Т., Пуура И. 1987. Кембрийско-ордовикские пограничные отложения разреза Юлгасе, Северная Эстония. — Изв. АН ЭССР. Геол., 36, 4, 154—165.
- Хейнсалу Х., Вийра В., Паалитс И. 1991. Пограничные кембро-ордовикские отложения разреза Сака II в Северо-Восточной Эстонии. — Изв. АН Эстонии. Геол., 40, 1, 8—15.

## ÜLEMKAMBRIUMI BIOSTRATIGRAAFIA EESTIS

Kaisa MENS, Viive VIIRA, Ivo PAALITS, Ivar PUURA

Ülemkambriumi läbilõigete kivimilise koostise ja ehituse järgi kuu- lub Eesti fatsiaalsetesse üleminekuvööndisse.

Ülemkambriumi biostratigraafiliseks liigestamiseks analüüsiti akri- tarhide, konodontide ja lingulaatide levikut mitmetes paljandites ning puursüdames. Saadud andmete põhjal on esitatud Eesti ülemkamb- riumi biotsonaalne liigestus, bioüksuste omavahelised suhted, samuti ka korrelatsioon trilobiitide tsoonidega.

## БИОСТРАТИГРАФИЯ ВЕРХНЕГО КЕМБРИЯ ЭСТОНИИ

Кайса МЕНС, Вийве ВИЙРА, Иво ПААЛИТС, Ивар ПУУРА

На основе анализа вертикального распределения акритарх, конодон- тов и лингулат выделены сообщества этих фоссилий и по ним рас- членен разрез верхнего кембрия Эстонии на биостратиграфические подразделения. По результатам сравнения родового и видового состав- вов этих сообществ с соответствующими материалами из разрезов, охарактеризованных трилобитами, выделенные биостратиграфические подразделения сопоставлены с трилобитовой зональной шкалой плат- формы (таблица).