

## Properties of 2-dimensional time-like ruled surfaces in the Minkowski space $\mathbb{R}_1^n$

Murat Tosun<sup>a</sup>, Ismail Aydemir<sup>b</sup>, and Nuri Kuruoglu<sup>b</sup>

<sup>a</sup> Department of Mathematics, Faculty of Arts and Science, Sakarya University, 54187 Sakarya, Turkey; tosun@sakarya.edu.tr

<sup>b</sup> Department of Mathematics, Faculty of Arts and Science, Ondokuz Mayıs University, Samsun, Turkey

Received 21 February 2005, in revised form 23 September 2005

**Abstract.** Some results, which are well known for the ruled surfaces in the Euclidean space  $\mathbb{R}^n$ , are generalized here to the case of  $\mathbb{R}_1^n$ . In particular, it is shown that a time-like ruled surface in  $\mathbb{R}_1^n$  is developable if and only if it has zero Gaussian curvature; moreover, it is then minimal if and only if it is totally geodesic.

**Key words:** ruled surfaces, Minkowski spaces.

### 1. INTRODUCTION

We shall assume throughout the paper that all manifolds, maps, vector fields, etc. are differentiable of class  $C^\infty$ . First of all, we give some properties of a general submanifold  $M$  of the Minkowski  $n$ -space  $\mathbb{R}_1^n$ . Suppose that  $\bar{D}$  is the Levi-Civita connection of  $\mathbb{R}_1^n$  and  $D$  is the Levi-Civita connection of  $M$ . Then, if  $X, Y$  are the vector fields of  $M$  and if  $V$  is the second fundamental tensor of  $M$ , we may decompose  $\bar{D}_X Y$  into a tangential and a normal component:

$$\bar{D}_X Y = D_X Y + V(X, Y). \quad (1)$$

Equation (1) is called Gauss equation [1]. If  $\xi$  is any normal vector field on  $M$ , we find the Weingarten equation by decomposing  $\bar{D}_X \xi$  into a tangential and a normal component:

$$\bar{D}_X \xi = -A_\xi(X) + D_X^\perp \xi, \quad (2)$$

where  $A_\xi$  determines at each point a self-adjoint linear map and  $D^\perp$  is a metric connection in the normal bundle  $\chi^\perp(M)$ . We use the same notation  $A_\xi$  for the linear map and the matrix of the linear map [2].

A normal vector field  $\xi$  is called parallel in the normal bundle  $\chi^\perp(M)$  if  $D_X^\perp \xi = 0$  for each vector field  $X$ . If  $\eta$  is a normal unit vector at the point  $p \in M$ , then

$$G(p, \eta) = \det A_\eta$$

is the Lipschitz–Killing curvature of  $M$  at  $p$  in the direction  $\eta$  [3].

Let  $V$  be the second fundamental tensor of  $M$ . If

$$V(X, X) = 0$$

for  $X$  in the tangent bundle  $\chi(M)$ , then  $X$  is called an asymptotic vector field on  $M$ . If

$$V(X, Y) = 0$$

for all  $X, Y \in \chi(M)$ , then  $M$  is totally geodesic [4].

Suppose that  $X, Y \in \chi(M)$ , while  $\xi \in \chi^\perp(M)$ . If the standard metric tensor of  $\mathbb{R}_1^n$  is denoted by  $\langle \cdot, \cdot \rangle$ , then we have

$$\langle \bar{D}_X Y, \xi \rangle = \langle V(X, Y), \xi \rangle$$

and

$$\langle \bar{D}_X Y, \xi \rangle = \langle A_\xi(X), Y \rangle.$$

From the above equations we obtain

$$\langle V(X, Y), \xi \rangle = \langle A_\xi(X), Y \rangle.$$

If  $\xi_1, \xi_2, \dots, \xi_{n-2}$  constitute an orthonormal base field of the normal bundle  $\chi^\perp(M)$ , then we set

$$\langle V(X, Y), \xi_j \rangle = V_j(X, Y)$$

or

$$V(X, Y) = \sum_{j=1}^{n-2} V_j(X, Y) \xi_j.$$

The mean curvature vector  $H$  of  $M$  at the point  $p$  is given by

$$H = \sum_{j=1}^{n-2} \frac{\text{tr} A_{\xi_j}}{2} \xi_j.$$

Here  $\|H\|$  is the mean curvature. If  $H = 0$  at each point  $p$  of  $M$ , then  $M$  is said to be minimal [5].

## 2. TWO-DIMENSIONAL TIME-LIKE RULED SURFACE IN $\mathbb{R}_1^n$

A time-like ruled surface  $M$  in  $\mathbb{R}_1^n$  is generated by time-like line  $l$  with unit direction time-like vector  $e(s)$  along a space-like curve  $\alpha$ . For this ruled surface

$$\psi(s, v) = \alpha(s) + ve(s)$$

is a parameterization. Throughout this paper,  $\alpha$  is supposed to be an orthogonal trajectory of the generators.

Let  $\{e, e_1\}$  be an orthonormal base field of  $\chi(M)$ , so that

$$\langle e, e \rangle = -1, \quad \langle e_1, e_1 \rangle = 1, \quad \langle e, e_1 \rangle = 0. \quad (3)$$

Let us denote the Levi-Civita connection of the Minkowski space  $\mathbb{R}_1^n$  by  $\bar{D}$ . Because the lines in  $\mathbb{R}_1^n$  are geodesics, we have

$$\bar{D}_e e = 0. \quad (4)$$

If we substitute this equation into Eq. (1), we get

$$V(e, e) = 0.$$

Considering Eq. (3), we can easily see that  $\bar{D}_e e_1 \perp e$  and  $\bar{D}_e e_1 \perp e_1$ . This implies  $\bar{D}_e e_1 \in \chi^\perp(M)$ . Therefore,

$$\bar{D}_e e_1 = V(e, e_1). \quad (5)$$

Let  $\{\xi_1, \xi_2, \dots, \xi_{n-2}\}$  be vector fields, which constitute an orthonormal base  $T_M^\perp(p)$ . Then  $\{e_1, e_2, \xi_1, \xi_2, \dots, \xi_{n-2}\}$  is a base of  $T_{\mathbb{R}_1^n}(p)$  at  $p \in \mathbb{R}_1^n$ . Together with (2) we can write

$$\begin{aligned} \bar{D}_e \xi_j &= a_{11}^j e + a_{12}^j e_1 + \sum_{i=1}^{n-2} b_{1i}^j \xi_i, \quad 1 \leq j \leq n-2, \\ \bar{D}_{e_1} \xi_j &= a_{21}^j e + a_{22}^j e_1 + \sum_{i=1}^{n-2} b_{2i}^j \xi_i, \quad 1 \leq j \leq n-2. \end{aligned} \quad (6)$$

Comparing Eqs. (6) with Eq. (4) leads us to

$$a_{21}^j = -a_{12}^j, \quad a_{11}^j = 0, \quad 1 \leq j \leq n-2.$$

Moreover, we find

$$A_{\xi_j} = \begin{bmatrix} 0 & a_{12}^j \\ -a_{12}^j & a_{22}^j \end{bmatrix}.$$

The matrix  $A_{\xi_j}$  corresponds to the shape operator of  $M$  and  $A_{\xi_j}$  is a symmetric matrix in the sense of Lorentz.

The Lipschitz–Killing curvature at  $p \in M$  in the direction of  $\xi_j$  is given by

$$G(p, \xi_j) = -(a_{12}^j)^2. \quad (7)$$

If we use Eqs. (6), we see

$$a_{12}^j = \langle \bar{D}_e \xi_j, e_1 \rangle = -\langle \xi_j, \bar{D}_e e_1 \rangle \quad (8)$$

and from (5) and with (8) we get

$$\bar{D}_e e_1 = V(e, e_1) = \sum_{j=1}^{n-2} \langle \xi_j, \bar{D}_e e_1 \rangle \xi_j = -\sum_{j=1}^{n-2} a_{12}^j \xi_j. \quad (9)$$

In addition, the Gaussian curvature of  $M$  denoted by  $G$  is expressed by (see [6])

$$G = -\langle \bar{D}_e e_1, \bar{D}_e e_1 \rangle.$$

With the elements of  $A_{\xi_j}$ , the Gaussian curvature of  $M$  is

$$G = -\sum_{j=1}^{n-2} (a_{12}^j)^2. \quad (10)$$

Hence, from Eqs. (7) and (10) we obtain

$$G(p) = \sum_{j=1}^{n-2} G(p, \xi_j), \quad p \in M. \quad (11)$$

Moreover, if the Lipschitz–Killing curvature  $G(p, \xi_j)$  is equal to zero at  $p \in M$  for each  $j$ ,  $1 \leq j \leq n-2$ , then Gaussian curvature  $G(p)$  will be zero. This shows that  $M$  is an intrinsically developable surface, i.e., locally isometric to open sets of Minkowski plane. Conversely, if  $M$  is intrinsically developable, then  $G(p, \xi_j)$  is equal to zero at  $p \in M$  for each  $j$ ,  $1 \leq j \leq n-2$ . Therefore, one may say that  $M$  is intrinsically developable if and only if the Lipschitz–Killing curvature is zero at each point [6].

In [6] it is shown that the mean curvature vector  $H$  of the time-like ruled surface  $M$  is

$$H = \frac{1}{2}V(e_1, e_1).$$

**Theorem 1.** *Let  $M$  be a 2-dimensional time-like ruled surface in  $\mathbb{R}_1^n$ . Then the generators of  $M$  are asymptotic and geodesic of  $M$ .*

*Proof.* Since the generators are the geodesics of  $\mathbb{R}_1^n$ , we write

$$\bar{D}_e e = 0.$$

If we set this into the Gauss equation, we find

$$D_e e + V(e, e) = 0 \text{ or } D_e e = -V(e, e).$$

Since  $D_e e \in \chi(M)$  and  $V(e, e) \in \chi^\perp(M)$  we reach  $D_e e = 0$  and  $V(e, e) = 0$ . This completes the proof of the theorem.

**Definition 1.** Let  $M$  be a 2-dimensional time-like ruled surface in  $\mathbb{R}_1^n$ . If the tangent planes of  $M$  are constant along the generators of  $M$ , then  $M$  is called developable [7].

**Theorem 2.** Let  $M$  be a 2-dimensional time-like ruled surface in  $\mathbb{R}_1^n$ . Then  $M$  is developable and minimal if and only if  $M$  is totally geodesic.

*Proof.* Assume that  $M$  is developable and minimal. If we have  $X = ae + be_1$  and  $Y = ce + de_1$  in  $\chi(M)$ , then

$$V(X, Y) = acV(e, e) + (ad + bc)V(e, e_1) + bdV(e_1, e_1).$$

Since the lines in  $\mathbb{R}_1^n$  are geodesic and  $M$  is minimal, we find that  $V(e, e) = V(e_1, e_1) = 0$ . Moreover,  $\bar{D}_e e_1$  is equal to zero since  $M$  is developable. From Eq. (5) we get

$$V(e, e_1) = 0.$$

Hence, we have  $V(X, Y) = 0$  for all  $X, Y \in \chi(M)$ . This means that  $M$  is totally geodesic.

Conversely, assume that  $V(X, Y) = 0$  for all  $X, Y \in \chi(M)$ . Therefore we have the relations

$$V(e, e) = 0, \quad V(e_1, e_1) = 0, \quad V(e, e_1) = 0.$$

By using these equations and Eq. (9) we find  $\bar{D}_e e_1 = 0$ . This shows that  $M$  is totally developable. Moreover,  $V(e_1, e_1) = 0$  implies that  $H = 0$ . This means that  $M$  is minimal.

### 3. SOME CHARACTERIZATIONS FOR 2-DIMENSIONAL TIME-LIKE RULED DEVELOPABLE SURFACES IN THE MINKOWSKI SPACE $\mathbb{R}_1^n$

Let  $\{e, e_1\}$  be an orthonormal basis of  $\chi(M)$ , as above, and  $\{\xi_1, \xi_2, \dots, \xi_{n-2}\}$  be an orthonormal basis of  $\chi^\perp(M)$ . We give covariant derivative equations of the orthonormal basis  $\{e_1, e_2, \xi_1, \xi_2, \dots, \xi_{n-2}\}$  of  $\chi(\mathbb{R}_1^n)$  as follows:

$$\begin{aligned}
\bar{D}_{e_1} e &= c_{11}e + c_{12}e_1 + c_{13}\xi_1 + \dots + c_{1n}\xi_{n-2}, \\
\bar{D}_{e_1} e_1 &= c_{21}e + c_{22}e_1 + c_{23}\xi_1 + \dots + c_{2n}\xi_{n-2}, \\
\bar{D}_{e_1} \xi_1 &= c_{31}e + c_{32}e_1 + c_{33}\xi_1 + \dots + c_{3n}\xi_{n-2}, \\
&\vdots \\
\bar{D}_{e_1} \xi_{n-2} &= c_{n1}e + c_{n2}e_1 + c_{n3}\xi_1 + \dots + c_{nn}\xi_{n-2}.
\end{aligned} \tag{12}$$

If we calculate the coefficient  $c_{st}$ ,  $1 \leq s, t \leq n$ , and write Eqs. (12) in the matrix form, we obtain

$$\begin{bmatrix} \bar{D}_{e_1} e \\ \bar{D}_{e_1} e_1 \\ \bar{D}_{e_1} \xi_1 \\ \vdots \\ \bar{D}_{e_1} \xi_{n-2} \end{bmatrix} = \begin{bmatrix} 0 & c_{12} & c_{13} & \cdots & c_{1n} \\ c_{12} & 0 & c_{23} & \cdots & c_{2n} \\ c_{13} & -c_{23} & 0 & \cdots & c_{3n} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ c_{1n} & -c_{2n} & -c_{3n} & \cdots & 0 \end{bmatrix} \begin{bmatrix} e \\ e_1 \\ \xi_1 \\ \vdots \\ \xi_{n-2} \end{bmatrix}. \tag{13}$$

By using Eq. (13) we can prove the following theorem.

**Theorem 3.** *Let  $M$  be a 2-dimensional time-like ruled surface in  $\mathbb{R}_1^n$ , and  $\{e, e_1\}$  be an orthonormal base field of the tangential bundle  $\chi(M)$ , as above. In this case, the following propositions are equivalent:*

- (i)  $M$  is developable,
- (ii) the Lipschitz–Killing curvature  $G(p, \xi_j)$ ,  $1 \leq j \leq n-2$ , is equal to zero,
- (iii) the Gaussian curvature  $G$  is equal to zero,
- (iv) in Eq. (13),  $c_{2k}$ ,  $3 \leq k \leq n$ , is equal to zero,
- (v)  $A_{\xi_j}^{\bar{D}_{e_1}}(e)$  is equal to zero,
- (vi)  $\bar{D}_{e_1} e_1$  is an element of  $\chi(M)$ .

*Proof.* (i)  $\Rightarrow$  (ii): Assume that  $M$  is developable, i.e.,  $\bar{D}_{e_1} e_1 = 0$ . Equation (7) says that the Lipschitz–Killing curvature at the point  $p$  in the direction of  $\xi_j$  is given by

$$G(p, \xi_j) = -(a_{12}^j(p))^2, \quad 1 \leq j \leq n-2. \tag{14}$$

Due to  $\bar{D}_{e_1} e_1 = 0$  and from Eq. (9)

$$\bar{D}_{e_1} e_1 = -\sum_{j=1}^{n-2} (a_{12}^j) \xi_j = 0. \tag{15}$$

Considering Eqs. (14) and (15) yields

$$G(p, \xi_j) = 0, \quad 1 \leq j \leq n-2.$$

(ii)  $\Rightarrow$  (iii): This follows directly from Eq. (11), as shown above.

(iii)  $\Rightarrow$  (iv): Assume that  $G = 0, \forall p \in M$ . From Eq. (10) we have  $a_{12}^j = 0, 1 \leq j \leq n-2$ . Since  $a_{21}^j = -a_{12}^j$  in (6), also  $a_{12}^j = 0$ . This means that  $\bar{D}_{e_1} \xi_j$  has no component in the direction  $e$ . Hence, we see that  $c_{2k} = 0, 3 \leq k \leq n$ , in Eqs. (12), due to (13).

(iv)  $\Rightarrow$  (v): Suppose that  $c_{2k} = 0, 3 \leq k \leq n$ , in Eqs. (12). This shows that  $\bar{D}_e \xi_j$  has no component in the direction  $e$ . Thus we have  $a_{12}^j = 0, 1 \leq j \leq n-2$ , in Eqs. (6).

Moreover, using Weingarten equation (2), we write

$$A_{\xi_j}(e) = 0, \quad 1 \leq j \leq n-2,$$

since  $a_{11}^j = -\langle \bar{D}_e \xi_j, e \rangle = \langle \xi_j, \bar{D}_e e \rangle = 0$ .

(v)  $\Rightarrow$  (vi): Let  $A_{\xi_j}(e)$  be equal to zero. Then, from Weingarten equation (2) we have  $a_{11}^j = 0, a_{12}^j = 0, 1 \leq j \leq n-2$ . Since  $\langle e, \xi_j \rangle = 0$  implies  $\langle \bar{D}_e e_1, \xi_j \rangle = \langle e, \bar{D}_{e_1} \xi_j \rangle = -a_{12}^j$ , we find

$$\langle \bar{D}_e e_1, \xi_j \rangle = 0.$$

From this equation we get

(vi)  $\Rightarrow$  (i): Let  $\bar{D}_e e_1$  be an element of  $\chi(M)$ . Then  $\langle \bar{D}_e e_1, \xi_j \rangle$  will be equal to  $-a_{12}^j, 1 \leq j \leq n-2$ , which is again equal to zero. On the other hand,  $\langle e_1, e_1 \rangle = 1$  implies that  $\langle \bar{D}_e e_1, e_1 \rangle = 0$  and  $\langle e_1, e \rangle = 0$  implies that  $\langle \bar{D}_e e_1, e \rangle = 0$ . Thus  $\bar{D}_e e_1 \in \chi^\perp(M)$ .

Using Eq. (9), we get that  $\bar{D}_e e_1 = 0$ , since  $a_{12}^j, 1 \leq j \leq n-2$ , is equal to zero. This means that the tangent planes of  $M$  are constant along the generator  $e$  of  $M$ , i.e.,  $M$  is developable. This finishes the proof.

## REFERENCES

1. Beem, J. K., Ehrlich, P. E. and Easley, K. L. *Global Lorentzian Geometry*, 2nd edition. Marcel Dekker, New York, 1996.
2. Chen, B. Y. *Geometry of Submanifolds*. Marcel Dekker, New York, 1973.
3. Houh, C. Surfaces with maximal Lipschitz–Killing curvature in the direction of mean curvature vector. *Proc. Amer. Math. Soc.*, 1972, **35**, 537–542.
4. Thas, C. Properties of ruled surfaces in the Euclidean space  $E^n$ . *Acad. Sinica*, 1978, **6**, 133–142.
5. Thas, C. *Een (lokale) studie van de  $(m+1)$ -dimensionale variëteiten, van de  $n$ -dimensionale euclidische ruimte  $R^n$  ( $n^3 2m + 1 \text{ en } m^3 1$ ), beschreven door een eendimensionale familie van  $m$ -dimensionale lineaire ruimten*. Palais Der Akademien-Herttogsstraat, I Brussel, 1974.
6. Tosun, M. and Aydemir, I. On 2-dimensional time-like ruled surfaces in the Minkowski space  $R_1^n$ . *Bull. Pure Appl. Sci.*, 1998, **17E**, 247–256.
7. O’Neill, B. *Semi-Riemannian Geometry*. Academic Press, New York, 1983.

# Ajasarnaste kahemõõtmeliste joonpindade omadusi Minkowski ruumis $\mathbb{R}_1^n$

Murat Tosun, Ismail Aydemir ja Nuri Kuruoglu

Mõned tulemused, mis on hästi tuntud joonpindade puhul eukleidilises ruumis  $\mathbb{R}^n$ , on üldistatud siin ruumi  $\mathbb{R}_1^n$  juhule. Nii on tõestatud, et ajasarnasel joonpinnal ruumis  $\mathbb{R}_1^n$  on puutujatasand piki iga moodustajat konstantne siis ja ainult siis, kui Gaussi kõverus on null; lisaks sellele on taoline joonpind minimaalne siis ja ainult siis, kui ta on täielikult geodeetiline.