

## On a class of Lorentzian para-Sasakian manifolds

Cengizhan Murathan<sup>a</sup>, Ahmet Yıldız<sup>b</sup>, Kadri Arslan<sup>a</sup>, and Uday Chand De<sup>c</sup>

<sup>a</sup> Department of Mathematics, Uludağ University, 16059 Bursa, Turkey; cengiz@uludag.edu.tr, arslan@uludag.edu.tr

<sup>b</sup> Department of Mathematics, Dumlupınar University, Kütahya, Turkey; ahmetyildiz@dumlupinar.edu.tr

<sup>c</sup> Department of Mathematics, Kalyani University, Kalyani, India; uc\_de@yahoo.com

Received 23 March 2006

**Abstract.** We classify Lorentzian para-Sasakian manifolds which satisfy  $P \cdot C = 0$ ,  $Z \cdot C = L_C Q(g, C)$ ,  $P \cdot Z - Z \cdot P = 0$ , and  $P \cdot Z + Z \cdot P = 0$ , where  $P$  is the  $v$ -Weyl projective tensor,  $Z$  is the concircular tensor, and  $C$  is the Weyl conformal curvature tensor.

**Key words:** contact metric manifold, Lorentzian para-Sasakian manifold, Sasakian manifold,  $v$ -Weyl projective tensor, concircular tensor.

### 1. INTRODUCTION

Matsumoto [1] introduced the notion of Lorentzian para-Sasakian ( $LP$ -Sasakian for short) manifold. Mihai and Rosca defined the same notion independently in [2]. This type of manifold is also discussed in [3,4].

Let  $M$  be an  $n$ -dimensional Riemannian manifold of class  $C^\infty$ . A  $v$ -projective symmetry is a projectable vector field  $X$  with the property in which every diffeomorphism  $\varphi$  of its one-parametric group is a projective map between leaves. In the theory of the projective transformations of connections the Weyl projective tensor plays an important role.

Recently, the authors of [5] studied the contact metric manifold  $M^n$  satisfying the curvature conditions  $Z(\xi, X) \cdot R = 0$  and  $R(\xi, X) \cdot Z = 0$ , where  $Z$  is the *concircular tensor* of  $M^n$  defined by

$$Z(X, Y)W = R(X, Y)W - \frac{\tau}{n(n-1)}R_0(X, Y)W, \quad (1)$$

where

$$R_0(X, Y)W = g(Y, W)X - g(X, W)Y,$$

$R$  and  $\tau$  are the *Riemannian–Christoffel curvature tensor* and the *scalar curvature* of  $M^n$ , respectively. They observed immediately from the form of the concircular curvature tensor that Riemannian manifolds with a vanishing concircular curvature tensor are of constant curvature. Thus one can think of the concircular curvature tensor as a measure of the failure of a Riemannian manifold to be of constant curvature.

In the theory of the projective transformations of connections the Weyl projective tensor plays an important role. The *v–Weyl projective tensor*  $P$  in a Riemannian manifold  $(M^n, g)$  is defined by [6]

$$P(X, Y)W = R(X, Y)W - \frac{1}{n-1}R_1(X, Y)W, \quad (2)$$

where

$$R_1(X, Y)W = S(Y, W)X - S(X, W)Y,$$

with  $S$  being the *Ricci tensor* of  $M$ .

In the present study we give a classification of the *LP-Sasakian manifold*  $M^n$  satisfying the curvature conditions  $P \cdot C = 0$ ,  $Z \cdot C = L_C Q(g, C)$ ,  $P \cdot Z - Z \cdot P = 0$ , and  $P \cdot Z + Z \cdot P = 0$ , where  $Z$  is the concircular tensor defined by (1),  $P$  is the *v–Weyl projective tensor*, and  $C$  is the *Weyl conformal curvature tensor* of  $M^n$ .

## 2. PRELIMINARIES

A differentiable manifold of dimension  $n$  is called an *LP-Sasakian manifold* [1,2] if it admits a  $(1, 1)$ -tensor field  $\phi$ , a contravariant vector field  $\xi$ , a covariant vector field  $\eta$ , and a Lorentzian metric  $g$  which satisfy

$$\eta(\xi) = -1, \quad (3)$$

$$\phi^2 = I + \eta \otimes \xi, \quad (4)$$

$$g(\phi X, \phi Y) = g(X, Y) + \eta(X)\eta(Y), \quad (5)$$

$$g(X, \xi) = \eta(X), \quad \nabla_X \xi = \phi X, \quad (6)$$

$$\Phi(X, Y) = g(X, \phi Y) = g(\phi X, Y) = \Phi(Y, X), \quad (7)$$

$$(\nabla_X \Phi)(Y, W) = g(Y, (\nabla_X \Phi)W) = (\nabla_X \Phi)(W, Y), \quad (8)$$

where  $\nabla$  is the covariant differentiation with respect to  $g$ . The Lorentzian metric  $g$  makes a timelike unit vector field, that is,  $g(\xi, \xi) = -1$ . The manifold  $M^n$  equipped with a Lorentzian almost paracontact structure  $(\phi, \xi, \eta, g)$  is said to be a *Lorentzian almost paracontact manifold* (see [1,3]).

If we replace in (3) and (4)  $\xi$  by  $-\xi$ , then the triple  $(\phi, \xi, \eta)$  is an almost paracontact structure on  $M^n$  defined by Sato [7]. The Lorentzian metric given by (6) stands analogous to the almost paracontact Riemannian metric for any almost paracontact manifold (see [7,8]).

A Lorentzian almost paracontact manifold  $M^n$  equipped with the structure  $(\phi, \xi, \eta, g)$  is called a *Lorentzian paracontact manifold* (see [1]) if

$$\Phi(X, Y) = \frac{1}{2} ((\nabla_X \eta)Y + (\nabla_Y \eta)X).$$

A Lorentzian almost paracontact manifold  $M^n$ , equipped with the structure  $(\phi, \xi, \eta, g)$ , is called an *LP-Sasakian manifold* (see [1]) if

$$(\nabla_X \phi)Y = g(\phi X, \phi Y)\xi + \eta(Y)\phi^2 X.$$

In an *LP-Sasakian manifold* the 1-form  $\eta$  is closed. In [1] it is also proved that if an  $n$ -dimensional Lorentzian manifold  $(M^n, g)$  admits a timelike unit vector field  $\xi$  such that the 1-form  $\eta$  associated to  $\xi$  is closed and satisfies

$$(\nabla_X \nabla_Y \eta)W = g(X, Y)\eta(W) + g(X, W)\eta(Y) + 2\eta(X)\eta(Y)\eta(W),$$

then  $M^n$  admits an *LP-Sasakian structure*.

Further, on such an *LP-Sasakian manifold*  $M^n$  with the structure  $(\phi, \xi, \eta, g)$  the following relations hold:

$$g(R(X, Y)W, \xi) = \eta(R(X, Y)W) = g(Y, W)\eta(X) - g(X, W)\eta(Y), \quad (9)$$

$$R(\xi, X)Y = g(X, Y)\xi - \eta(Y)X, \quad (10)$$

$$R(X, Y)\xi = \eta(Y)X - \eta(X)Y, \quad (11)$$

$$R(\xi, X)\xi = X + \eta(X)\xi, \quad (12)$$

$$S(X, \xi) = (n - 1)\eta(X), \quad (13)$$

$$S(\phi X, \phi Y) = S(X, Y) + (n - 1)\eta(X)\eta(Y) \quad (14)$$

for any vector fields  $X, Y$  (see [1,2]), where  $S$  is the Ricci curvature and  $Q$  is the Ricci operator given by  $S(X, Y) = g(QX, Y)$ .

An *LP-Sasakian manifold*  $M^n$  is said to be  $\eta$ -Einstein if its Ricci tensor  $S$  is of the form

$$S(X, Y) = ag(X, Y) + b\eta(X)\eta(Y) \quad (15)$$

for any vector fields  $X, Y$ , where  $a, b$  are functions on  $M^n$  (see [9,10]).

Next we define endomorphisms  $R(X, Y)$  and  $X \wedge_A Y$  of  $\chi(M)$  by

$$R(X, Y)W = \nabla_X \nabla_Y W - \nabla_Y \nabla_X W - \nabla_{[X, Y]}W, \quad (16)$$

$$(X \wedge_A Y)W = A(Y, W)X - A(X, W)Y, \quad (17)$$

respectively, where  $X, Y, W \in \chi(M)$  and  $A$  is the symmetric  $(0, 2)$ -tensor.

For a  $(0, k)$ -tensor field  $T$ ,  $k \geq 1$ , on  $(M, g)$  we define  $P \cdot T$ ,  $Z \cdot T$ , and  $Q(g, T)$  by

$$\begin{aligned} (P(X, Y) \cdot T)(X_1, \dots, X_k) &= -T(P(X, Y)X_1, X_2, \dots, X_k) \\ &\quad -T(X_1, P(X, Y)X_2, \dots, X_k) \\ &\quad - \dots - T(X_1, X_2, \dots, P(X, Y)X_k), \end{aligned} \quad (18)$$

$$\begin{aligned} (Z(X, Y) \cdot T)(X_1, \dots, X_k) &= -T(Z(X, Y)X_1, X_2, \dots, X_k) \\ &\quad -T(X_1, Z(X, Y)X_2, \dots, X_k) \\ &\quad - \dots - T(X_1, X_2, \dots, Z(X, Y)X_k), \end{aligned} \quad (19)$$

$$\begin{aligned} Q(g, T)(X_1, \dots, X_k; X, Y) &= -T((X \wedge Y)X_1, X_2, \dots, X_k) \\ &\quad -T(X_1, (X \wedge Y)X_2, \dots, X_k) \\ &\quad - \dots - T(X_1, X_2, \dots, (X \wedge Y)X_k), \end{aligned} \quad (20)$$

respectively [11].

By definition the Weyl conformal curvature tensor  $C$  is given by

$$\begin{aligned} C(X, Y)Z &= R(X, Y)Z - \frac{1}{n-2} \left[ \begin{array}{l} g(Y, Z)QX - g(X, Z)QY \\ +S(Y, Z)X - S(X, Z)Y \end{array} \right] \\ &\quad + \frac{\tau}{(n-1)(n-2)} [g(Y, Z)X - g(X, Z)Y], \end{aligned} \quad (21)$$

where  $Q$  denotes the Ricci operator, i.e.,  $S(X, Y) = g(QX, Y)$  and  $\tau$  is scalar curvature [9]. The Weyl conformal curvature tensor  $C$  is defined by  $C(X, Y, Z, W) = g(C(X, Y)Z, W)$ . If  $C = 0$ ,  $n \geq 4$ , then  $M$  is called conformally flat.

### 3. MAIN RESULTS

In the present section we consider the  $LP$ -Sasakian manifold  $M^n$  satisfying the curvature conditions  $P \cdot C = 0$ ,  $Z \cdot C = L_C Q(g, C)$ ,  $P \cdot Z - Z \cdot P = 0$ , and  $P \cdot Z + Z \cdot P = 0$ .

First we give the following proposition.

**Proposition 1.** *Let  $M$  be an  $n$ -dimensional ( $n > 3$ )  $LP$ -Sasakian manifold. If the condition  $P \cdot C = 0$  holds on  $M$ , then*

$$\begin{aligned} S^2(X, U) &= \left[ \frac{\tau}{n-1} - (n-1)^2 - 1 \right] S(X, U) \\ &\quad + (n-1)[\tau - (n-1)]g(X, U) \\ &\quad + n[\tau - n(n-1)]\eta(X)\eta(U) \end{aligned}$$

is satisfied on  $M$ , where  $S^2(X, U) = S(QX, U)$ .

*Proof.* Assume that  $M$  is an  $n$ -dimensional,  $n > 3$ ,  $LP$ -Sasakian manifold satisfying the condition  $P \cdot C = 0$ . From (18) we have

$$\begin{aligned} (P(V, X) \cdot C)(Y, U)W &= P(V, X)C(Y, U)W \\ &\quad - C(P(V, X)Y, U)W - C(Y, P(V, X)U)W \\ &\quad - C(Y, U)P(V, X)W = 0, \end{aligned} \quad (22)$$

where  $X, Y, U, V, W \in \chi(M)$ . Taking  $V = \xi$  in (22), we have

$$\begin{aligned} (P(\xi, X) \cdot C)(Y, U)W &= P(\xi, X)C(Y, U)W \\ &\quad - C(P(\xi, X)Y, U)W - C(Y, P(\xi, X)U)W \\ &\quad - C(Y, U)P(\xi, X)W = 0. \end{aligned} \quad (23)$$

Furthermore, substituting (2), (9), (13), (21) into (23) and multiplying with  $\xi$ , we get

$$\begin{aligned} &-g(X, C(Y, U)W) - n\eta(C(Y, U)W)\eta(X) - g(X, Y)\eta(C(\xi, U)W) \\ &+ n\eta(Y)\eta(C(X, U)W) - g(X, U)\eta(C(Y, \xi)W) \\ &+ n\eta(U)\eta(C(Y, X)W) + n\eta(W)\eta(C(Y, U)X) \\ &+ \frac{1}{n-1}\{S(X, C(Y, U)W) + S(X, Y)\eta(C(\xi, U)W) \\ &+ S(X, U)\eta(C(Y, \xi)W)\} = 0. \end{aligned} \quad (24)$$

Thus, replacing  $W$  with  $\xi$  in (24), we have

$$-g(X, C(Y, U)\xi) - n\eta(C(Y, U)X) + \frac{1}{n-1}S(X, C(Y, U)\xi) = 0. \quad (25)$$

Again, taking  $Y = \xi$  in (25) and after some calculations, since  $n > 3$ , we get

$$\begin{aligned} S^2(X, U) &= \left[ \frac{\tau}{n-1} - (n-1)^2 - 1 \right] S(X, U) \\ &\quad + (n-1)[\tau - (n-1)]g(X, U) \\ &\quad + n[\tau - n(n-1)]\eta(X)\eta(U). \end{aligned}$$

Our theorem is thus proved.  $\square$

**Theorem 2.** Let  $M$  be an  $n$ -dimensional ( $n > 3$ )  $LP$ -Sasakian manifold. If the condition  $Z \cdot C = L_C Q(g, C)$  holds on  $M$ , then either  $M$  is conformally flat or  $L_C = \frac{\tau}{n(n-1)} - 1$ .

*Proof.* Let  $M^n$  be an  $LP$ -Sasakian manifold. So we have

$$(Z(V, X) \cdot C)(Y, U)W = L_C Q(g, C)(Y, U, W; V, X).$$

Then from (19) and (20) we can write

$$\begin{aligned} & Z(V, X)C(Y, U)W - C(Z(V, X)Y, U)W - C(Y, Z(V, X)U)W \\ & \quad - C(Y, U)Z(V, X)W \\ & = L_C[(V \wedge X)C(Y, U)W - C((V \wedge X)Y, U)W \\ & \quad - C(Y, (V \wedge X)U)W - C(Y, U)(V \wedge X)W]. \end{aligned} \quad (26)$$

Therefore, replacing  $V$  with  $\xi$  in (26), we have

$$\begin{aligned} & Z(\xi, X)C(Y, U)W - C(Z(\xi, X)Y, U)W - C(Y, Z(\xi, X)U)W \\ & \quad - C(Y, U)Z(\xi, X)W \\ & = L_C[(\xi \wedge X)C(Y, U)W - C((\xi \wedge X)Y, U)W \\ & \quad - C(Y, (\xi \wedge X)U)W - C(Y, U)(\xi \wedge X)W]. \end{aligned} \quad (27)$$

Using (20), (9) and taking the inner product of (27) with  $\xi$ , we get

$$\begin{aligned} & \left[1 - \frac{\tau}{n(n-1)} - L_C\right] [-g(X, C(Y, U)W) - \eta(C(Y, U)W)\eta(X) \\ & \quad - g(X, Y)\eta(C(\xi, U)W) + \eta(Y)\eta(C(X, U)W) \\ & \quad - g(X, U)\eta(C(Y, \xi)W) + \eta(U)\eta(C(Y, X)W) + \eta(W)\eta(C(Y, U)X)] = 0. \end{aligned} \quad (28)$$

Putting  $X = Y$  in (28), we have

$$\begin{aligned} & \left[1 - \frac{\tau}{n(n-1)} - L_C\right] [-g(Y, C(Y, U)W) + \eta(W)\eta(C(Y, U)Y) \\ & \quad - g(Y, Y)\eta(C(\xi, U)W) - g(Y, U)\eta(C(Y, \xi)W)] = 0. \end{aligned} \quad (29)$$

A contraction of (29) with respect to  $Y$  gives us

$$\left[1 - \frac{\tau}{n(n-1)} - L_C\right] \eta(C(\xi, U)W) = 0. \quad (30)$$

If  $L_C \neq 1 - \frac{\tau}{n(n-1)}$ , then Eq. (30) is reduced to

$$\eta(C(\xi, U)W) = 0, \quad (31)$$

which gives

$$S(U, W) = \left(\frac{\tau}{(n-1)} - 1\right) g(U, W) + \left(\frac{\tau}{(n-1)} - n\right) \eta(U)\eta(W). \quad (32)$$

Therefore,  $M$  is a  $\eta$ -Einstein manifold. So, using (31) and (32), we have Eq. (28) in the form

$$C(Y, U, W, X) = 0,$$

which means that  $M$  is conformally flat.

If  $L_C \neq 0$  and  $\eta(C(\xi, U)W) \neq 0$ , then  $1 - \frac{\tau}{n(n-1)} - L_C = 0$ , which gives  $L_C = 1 - \frac{\tau}{n(n-1)}$ . This completes the proof of the theorem.  $\square$

**Corollary 3.** Every  $n$ -dimensional ( $n > 3$ ) nonconformally flat LP-Sasakian manifold satisfies  $Z \cdot C = (1 - \frac{\tau}{n(n-1)})Q(g, C)$ .

**Theorem 4.** Let  $M$  be an  $n$ -dimensional ( $n > 3$ ) LP-Sasakian manifold.  $M$  satisfies the condition

$$P \cdot Z - Z \cdot P = 0$$

if and only if  $M$  is a  $\eta$ -Einstein manifold.

*Proof.* Let  $M$  satisfy the condition  $P \cdot Z - Z \cdot P = 0$ . Then we can write

$$\begin{aligned} & P(V, X) \cdot Z(Y, U)W - Z(V, X) \cdot P(Y, U)W \\ &= \frac{1}{n-1} [R(V, X)R_1(Y, U)W - R_1(V, X)R(Y, U)W] \\ & \quad + \frac{\tau}{n(n-1)^2} [R_1(V, X)R_0(Y, U)W - R_0(V, X)R_1(Y, U)W] \\ & \quad + \frac{\tau}{n(n-1)} [R_0(V, X)R(Y, U)W - R(V, X)R_0(Y, U)W] = 0. \quad (33) \end{aligned}$$

Therefore, replacing  $V$  with  $\xi$  in (33), we have

$$\begin{aligned} & P(\xi, X) \cdot Z(Y, U)W - Z(\xi, X) \cdot P(Y, U)W \\ &= \frac{1}{n-1} [R(\xi, X)R_1(Y, U)W - R_1(\xi, X)R(Y, U)W] \\ & \quad + \frac{\tau}{n(n-1)^2} [R_1(\xi, X)R_0(Y, U)W - R_0(\xi, X)R_1(Y, U)W] \\ & \quad + \frac{\tau}{n(n-1)} [R_0(\xi, X)R(Y, U)W - R(\xi, X)R_0(Y, U)W] = 0. \quad (34) \end{aligned}$$

Using (10), (13), we get

$$\begin{aligned} & \frac{1}{n-1} [S(U, W)g(X, Y)\xi - S(U, W)\eta(Y)X - g(X, U)S(Y, W)\xi \\ & \quad + S(Y, W)\eta(U)X - S(X, R(Y, U)W)\xi + (n-1)g(U, W)\eta(Y)X \\ & \quad - (n-1)g(Y, W)\eta(U)X] \\ & \quad + \frac{\tau}{n(n-1)^2} [g(U, W)g(X, Y)\xi - g(U, W)\eta(Y)X - g(Y, W)g(X, U)\xi \\ & \quad + g(Y, W)\eta(U)X - S(U, W)g(X, Y)\xi + S(U, W)\eta(Y)X \\ & \quad + S(Y, W)g(X, U)\xi - S(Y, W)\eta(U)X] \\ & \quad + \frac{\tau}{n(n-1)} [g(X, R(Y, U)W)\xi + g(Y, W)\eta(U)X - g(U, W)g(X, Y)\xi \\ & \quad + g(Y, W)g(X, U)\xi - g(Y, W)\eta(U)X] = 0. \quad (35) \end{aligned}$$

Again, taking  $U = \xi$  in (35), we get

$$\begin{aligned} & \frac{1}{n-1} [(n-1)g(X, Y)\eta(W)\xi - S(Y, W)\eta(X)\xi - S(Y, W)X \\ & + (n-1)g(Y, W)\eta(X)\xi - S(X, Y)\eta(W)\xi + (n-1)g(Y, W)X] \\ & + \frac{\tau}{n(n-1)^2} [g(X, Y)\eta(W)\xi - \eta(W)\eta(Y)X - g(Y, W)\eta(X)\xi - g(Y, W)X \\ & - (n-1)g(X, Y)\eta(W)\xi + (n-1)\eta(W)\eta(Y)X \\ & - S(Y, W)\eta(X)\xi + S(Y, W)X] = 0. \end{aligned} \quad (36)$$

Taking the inner product of (36) with  $\xi$ , we find

$$\begin{aligned} & \frac{1}{n-1} [S(X, Y)\eta(W) - (n-1)g(X, Y)\eta(W)] \\ & + \frac{\tau(n-2)}{n(n-1)^2} [g(X, Y)\eta(W) + \eta(X)\eta(Y)\eta(W)] = 0. \end{aligned} \quad (37)$$

Again, taking  $W = \xi$  and using (3) in (37), we get

$$\begin{aligned} S(X, Y) &= \left[ (n-1) - \frac{(n-2)}{n(n-1)}\tau \right] g(X, Y) \\ & - \left[ \frac{(n-2)}{n(n-1)}\tau \right] \eta(X)\eta(Y). \end{aligned} \quad (38)$$

So,  $M$  is a  $\eta$ -Einstein manifold.

Conversely, if  $M^n$  is a  $\eta$ -Einstein manifold, then it is easy to show that  $P \cdot Z - Z \cdot P = 0$ . Our theorem is thus proved.  $\square$

**Theorem 5.** *Let  $M$  be an  $n$ -dimensional ( $n > 3$ ) LP-Sasakian manifold.  $M$  satisfies the condition*

$$P \cdot Z + Z \cdot P = 0$$

*if and only if  $M$  is an Einstein manifold.*

*Proof.* Let  $M$  satisfy the condition  $P \cdot Z + Z \cdot P = 0$ . Then, from (33) and (34), we can write

$$\begin{aligned} & 2R(\xi, X)R(Y, U)W \\ & - \frac{1}{n-1} [R(\xi, X)R_1(Y, U)W + R_1(\xi, X)R(Y, U)W] \\ & + \frac{\tau}{n(n-1)^2} [R_1(\xi, X)R_0(Y, U)W + R_0(\xi, X)R_1(Y, U)W] \\ & - \frac{\tau}{n(n-1)} [R_0(\xi, X)R(Y, U)W + R(\xi, X)R_0(Y, U)W] = 0. \end{aligned} \quad (39)$$



Using (6), (10), and (13) in (39), we have

$$\begin{aligned}
& 2[g(X, R(Y, U)W)\xi - g(U, W)\eta(Y)X + g(Y, W)\eta(U)X] \\
& - \frac{1}{n-1}[S(U, W)g(X, Y)\xi - S(U, W)\eta(Y)X - S(Y, W)g(X, U)\xi \\
& + S(Y, W)\eta(U)X + S(X, R(Y, U)W)\xi - (n-1)g(U, W)\eta(Y)X \\
& + (n-1)g(Y, W)\eta(U)X] \\
& + \frac{\tau}{n(n-1)^2}[g(U, W)S(X, Y)\xi - (n-1)g(U, W)\eta(Y)X \\
& - g(Y, W)S(X, U)\xi + (n-1)g(Y, W)\eta(U)X + S(U, W)g(X, Y)\xi \\
& - S(U, W)\eta(Y)X - S(Y, W)g(X, U)\xi + S(Y, W)\eta(U)X] \\
& - \frac{\tau}{n(n-1)}[g(X, R(Y, U)W)\xi - 2g(U, W)\eta(Y)X + 2g(Y, W)\eta(U)X \\
& + g(U, W)g(X, Y)\xi - g(Y, W)g(X, U)\xi] = 0. \tag{40}
\end{aligned}$$

Replacing  $Y$  with  $\xi$  and using (3) in (40), we have

$$\begin{aligned}
& 2[g(X, R(\xi, U)W)\xi + g(U, W)X + \eta(W)\eta(U)X] \\
& - \frac{1}{n-1}[S(U, W)\eta(X)\xi + S(U, W)X - (n-1)g(X, U)\eta(W)\xi \\
& + 2(n-1)\eta(W)\eta(U)X + S(X, R(\xi, U)W)\xi + (n-1)g(U, W)X] \\
& + \frac{\tau}{n(n-1)^2}[(n-1)g(U, W)\eta(X)\xi + (n-1)g(U, W)X \\
& - S(X, U)\eta(W)\xi + (n-1)\eta(W)\eta(U)X + S(U, W)\eta(X)\xi \\
& + S(U, W)X - (n-1)g(X, U)\eta(W)\xi + (n-1)\eta(W)\eta(U)X] \\
& - \frac{\tau}{n(n-1)}[g(X, R(\xi, U)W)\xi + 2g(U, W)X + 2\eta(W)\eta(U)X \\
& + g(U, W)\eta(X)\xi - g(X, U)\eta(W)\xi] = 0. \tag{41}
\end{aligned}$$

Taking the inner product of (41) with  $\xi$  and using (7), (10), we get

$$\begin{aligned}
& \left[2 - \frac{2\tau}{n(n-1)}\right] [g(X, U)\eta(W) + \eta(X)\eta(U)\eta(W)] \\
& + \left[\frac{\tau}{n(n-1)^2} - \frac{1}{n-1}\right] [(n-1)g(X, U)\eta(W) + 2(n-1)\eta(X)\eta(U)\eta(W) \\
& + S(X, U)\eta(W)] = 0. \tag{42}
\end{aligned}$$

Again, taking  $W = \xi$  and using (3) in (42), we get

$$\begin{aligned}
& \left[\frac{2\tau}{n(n-1)} - 2\right] [g(X, U) + \eta(X)\eta(U)] \\
& - \left[\frac{\tau}{n(n-1)^2} - \frac{1}{n-1}\right] [(n-1)g(X, U) \\
& + 2(n-1)\eta(X)\eta(U) + S(X, U)] = 0. \tag{43}
\end{aligned}$$

Thus, from (43), we have

$$S(X, U) = (n - 1)g(X, U).$$

So,  $M^n$  is an Einstein manifold.

Conversely, if  $M^n$  is an Einstein manifold, then it is easy to show that  $P \cdot Z + Z \cdot P = 0$ . Our theorem is thus proved.  $\square$

### ACKNOWLEDGEMENT

This study was supported by the Dumlupınar University research foundation (project No. 2004-9).

### REFERENCES

1. Matsumoto, K. On Lorentzian paracontact manifolds. *Bull. of Yamagata Univ. Nat. Sci.*, 1989, **12**, 151–156.
2. Mihai, I. and Rosca, R. *On Lorentzian P-Sasakian Manifolds, Classical Analysis*. World Scientific, Singapore, 1992, 155–169.
3. Matsumoto, K. and Mihai, I. On a certain transformation in a Lorentzian para-Sasakian manifold. *Tensor, N. S.*, 1988, **47**, 189–197.
4. Tripathi, M. M. and De, U. C. Lorentzian almost paracontact manifolds and their submanifolds. *J. Korea Soc. Math. Educ. Ser. B: Pure Appl. Math.*, 2001, **8**, 101–105.
5. Blair, D. E., Kim, J. S. and Tripathi, M. M. On the concircular curvature tensor of a contact metric manifold. *J. Korean Math. Soc.*, 2005, **42**, 883–892.
6. Tığaerü, C.  $v$ -projective symmetries of fibered manifolds. *Arch. Math.*, 1998, **34**, 347–352.
7. Sato, I. On a structure similar to almost contact structures. *Tensor, N. S.*, 1976, **30**, 219–224.
8. Sato, I. On a structure similar to almost contact structures II. *Tensor, N. S.*, 1977, **31**, 199–205.
9. Yano, K. and Kon, M. *Structures on Manifolds*. Series in Pure Mathematics, Vol. 3, 1984. World Scientific, Singapore.
10. Blair, D. E. *Contact Manifolds in Riemannian Geometry*. Lecture Notes in Mathematics, Vol. 509, 1976, Springer-Verlag, Berlin.
11. Deszcz, R. On pseudosymmetric spaces. *Bull. Soc. Math. Belg.*, 1990, **49**, 134–145.

## Ühest Lorentzi para-Sasaki muutkondade klassist

Cengizhan Murathan, Ahmet Yıldız, Kadri Arslan ja Uday Chand De

On käsitletud Lorentzi para-Sasaki muutkondi, mille puhul  $P \cdot C = 0$ ,  $Z \cdot C = L_C Q(g, C)$ ,  $P \cdot Z - Z \cdot P = 0$  või  $P \cdot Z + Z \cdot P = 0$ , kus  $C$  on Weyli konformse kõveruse tensor,  $P$  on  $v$ -Weyli projektiivne tensor ja  $Z$  on kantsirkulaartensor.