

EESTI NSV TEADUSTE AKADEEMIA TOIMETISED. 25. KOIDE
FÜSIKA * MATEMAATIKA. 1976, NR. 1

ИЗВЕСТИЯ АКАДЕМИИ НАУК ЭСТОНСКОЙ ССР. ТОМ 25
ФИЗИКА * МАТЕМАТИКА. 1976, № 1

УДК 539.12

M. KOIV, R.-K. LOIDE

A NOTE ON SUPERFIELD ALGEBRAS

M. KOIV, R.-K. LOIDE. MÄRKUS SUPervalJADE ALGEBRATE KONTA

M. КИЙВ, Р.-К. ЛОЙДЕ. ЗАМЕЧАНИЕ ОБ АЛГЕБРАХ СУПЕРПОЛЕЙ

Supersymmetry introduced by Wess and Zumino [1,2] has recently gained great interest [3-5]. In the case of constant parameters this symmetry was presented by Salam and Strathdee in the form of graded Lie algebras studied from a purely mathematical point of view in [6,7].

In this note we deal with the Wess-Zumino type of superalgebras where the number of supergauge generators is arbitrary, and give the general form of the structure constants. We consider an algebra where the generators P^μ and $M^{\mu\nu}$ of the Poincaré group and supergauge generators S_α satisfy the relations

$$\begin{aligned} [P^\mu, P^\nu] &= 0 \\ [M^{\mu\nu}, P^\rho] &= c_{\kappa}^{\mu\nu, \rho} P^\kappa \\ [M^{\mu\nu}, M^{\rho\sigma}] &= c_{\kappa\lambda}^{\mu\nu, \rho\sigma} M^{\kappa\lambda} \\ [S_\alpha, P^\mu] &= 0 \\ [S_\alpha, M^{\mu\nu}] &= B_{\alpha\beta}^{\mu\nu} S_\beta \\ \{S_\alpha, S_\beta\} &= A_{\kappa, \alpha\beta} P^\kappa \end{aligned} \quad (1)$$

Here

$$\begin{aligned} c_{\kappa\lambda}^{\mu\nu, \rho\sigma} &= \frac{1}{2} [g^{\mu\sigma} (g_\kappa^\nu g_\lambda^\rho - g_\kappa^\rho g_\lambda^\nu) + g^{\nu\rho} (g_\kappa^\mu g_\lambda^\sigma - g_\kappa^\sigma g_\lambda^\mu) - \\ &\quad - g^{\mu\rho} (g_\kappa^\nu g_\lambda^\sigma - g_\kappa^\sigma g_\lambda^\nu) - g^{\nu\sigma} (g_\kappa^\mu g_\lambda^\rho - g_\kappa^\rho g_\lambda^\mu)], \end{aligned} \quad (2)$$

$$c_{\kappa}^{\mu\nu, \rho} = g^{\nu\rho} g_\kappa^\mu - g^{\mu\rho} g_\kappa^\nu \quad (3)$$

and $g^{00} = -g^{11} = -g^{22} = -g^{33} = 1$. In this note $\kappa, \lambda, \mu, \nu, \rho, \sigma = 0, 1, 2, 3$ and $\alpha, \beta, \gamma, \delta = 1, 2, \dots, n$, summation over the repeating indices is

presumed. Proceeding from the Jacobi identities, we shall show that $B_{\alpha\beta}^{\mu\nu}$ are the matrix elements of the generators of some representation of the Lorentz group and $A_{\mu,\alpha\beta}$ are connected with the β -matrices of an invariant equation $(p_\mu\beta^\mu - m)\psi(p) = 0$. The idea is to reduce the conditions we get from the Jacobi identities to some well-known commutation relations. Further we treat the structure constants $B_{\alpha\beta}^{\mu\nu}$ and $A_{\mu,\alpha\beta}$ as the matrix elements of matrices $B^{\mu\nu}$ and A_μ .

Starting with the Jacobi identities it is easy to be convinced that we must look through the identities with generators S_α , $M^{\mu\nu}$, $M^{\rho\sigma}$ and S_β , $M^{\mu\nu}$, S_α .

$$[S_\alpha, [M^{\mu\nu}, M^{\rho\sigma}]] + [M^{\rho\sigma}, [S_\alpha, M^{\mu\nu}]] + [M^{\mu\nu}, [M^{\rho\sigma}, S_\alpha]] = 0.$$

Using the relations (1), we obtain

$$c_{\kappa\lambda}^{\mu\nu,\rho\sigma} B_{\alpha\gamma}^{\kappa\lambda} - B_{\alpha\beta}^{\mu\nu} B_{\beta\gamma}^{\rho\sigma} + B_{\alpha\beta}^{\rho\sigma} B_{\beta\gamma}^{\mu\nu} = 0.$$

That gives for matrices $B^{\mu\nu}$

$$[B^{\mu\nu}, B^{\rho\sigma}] = c_{\kappa\lambda}^{\mu\nu,\rho\sigma} B^{\kappa\lambda}. \quad (4)$$

The last relations are the commutation relations of the Lorentz group. Therefore the matrices $B^{\mu\nu}$ are the generators of some arbitrary representation of the Lorentz group, and the structure constants $B_{\alpha\beta}^{\mu\nu}$ are its matrix elements. The number of supergauge generators is equal to the dimension of the $B^{\mu\nu}$ representation.

The second Jacobi identity

$$-\{S_\beta, [M^{\mu\nu}, S_\alpha]\} + \{S_\alpha, [S_\beta, M^{\mu\nu}]\} + [M^{\mu\nu}, \{S_\alpha, S_\beta\}] = 0$$

gives the relations

$$B_{\alpha\gamma}^{\mu\nu} A_{\kappa,\beta\gamma} + B_{\beta\gamma}^{\mu\nu} A_{\kappa,\alpha\gamma} + c_{\kappa}^{\mu\nu,\lambda} A_{\lambda,\alpha\beta} = 0.$$

For matrices $B^{\mu\nu}$ and A_μ

$$B^{\mu\nu} A_\kappa + A_\kappa \bar{B}^{\mu\nu} = -c_\kappa^{\mu\nu,\lambda} A_\lambda. \quad (5)$$

Here $\bar{B}^{\mu\nu}$ is $B^{\mu\nu}$ transposed and we have used the fact that $\bar{A}_\mu = A_\mu$.

Now we suppose that there exists a matrix C which satisfies

$$C \bar{B}^{\mu\nu} C^{-1} = -B^{\mu\nu}. \quad (6)$$

Then we can write (5) in the form

$$[B^{\mu\nu}, A_\kappa C^{-1}] = -c_\kappa^{\mu\nu,\lambda} A_\lambda C^{-1}.$$

Denoting

$$\beta_\kappa = A_\kappa C^{-1} \quad (7)$$

we obtain

$$[B^{\mu\nu}, \beta_\kappa] = -c_\kappa^{\mu\nu,\lambda} \beta_\lambda \quad (8)$$

The last relations are the invariance conditions for the first-order wave equation corresponding to the $B^{\mu\nu}$ representation. Therefore the matrices A_κ are connected with the β -matrices of an invariant equation.

From the relation (7)

$$A_\kappa = \beta_\kappa C \quad (9)$$

and therefore

$$\{S_\alpha, S_\beta\} = (\beta_\kappa C)_{\alpha\beta} P^\kappa$$

As regards matrix C which satisfies (6), it should be mentioned that such a matrix always exists because it exists in the case of the spinor representations $(1/2, 0)$ and $(0, 1/2)$. To show this, we write the generators $B^{\mu\nu}$ in the form: $B^{0k} = \mp \frac{1}{2} \sigma^k$, $B^{kl} = \frac{i}{2} \epsilon_{lm}^k \sigma^m$ ($k, l, m = 1, 2, 3$; $\epsilon^{123} = +1$) where the minus sign corresponds to the representation $(1/2, 0)$, and the plus sign to $(0, 1/2)$ and $\sigma^1, \sigma^2, \sigma^3$ are the Pauli matrices. Now C must satisfy $C \sigma^k C^{-1} = -\sigma^k$. In the case of the usual representation of the Pauli matrices one can take $C = \pm \sigma^2$. As the other representations are obtainable from the direct products of spinor representations, matrix C also exists. In the case of the Dirac bispinor $(1/2, 0) \oplus (0, 1/2)$ C is the charge conjugation operator.

The matrices β_* and C must be chosen to satisfy $\beta_* C = \beta_* C$.

Some conclusions:

1) In the case of the irreducible representation of $B^{\mu\nu}$ the matrices $A_* = 0$ because there exists no first-order invariant equation. Therefore in this case $\{S_\alpha, S_\beta\} = 0$. From (5) it is easy to see that $A_* = 0$ is always one of the possible solutions.

2) The Wess-Zumino algebra is the only one which has four supergauge generators S_α and $\{S_\alpha, S_\beta\} \neq 0$. The vector representation $(1/2, 1/2)$ has also four generators S_α but due to the previously mentioned we get $\{S_\alpha, S_\beta\} = 0$.

3) The next possible superalgebra we get after the Wess-Zumino algebra is the algebra connected with the Kemmer-Duffin matrices β^μ for spin 0 (Wess-Zumino algebra is connected with the Dirac matrices γ^μ). In this case there are five supergenerators S_α , $B^{\mu\nu} = [\beta^\mu, \beta^\nu]$ and C is the space reflection operator which must be chosen to satisfy $A_* = A_*$. The superfield connected with the Kemmer-Duffin matrices was recently studied by Ainsaar [8].

We are indebted to A. Ainsaar and R. Saar for several enlightening discussions.

REFERENCES

1. Wess J., Zumino B., Phys. Lett., **B49**, 52 (1974).
2. Wess J., Zumino B., Nucl. Phys., **B70**, 39 (1974).
3. Salam A., Strathdee J., Nucl. Phys., **B76**, 477 (1974).
4. Salam A., Strathdee J., Unitary representations of supergauge symmetries, Preprint IC/74/16, Trieste, 1974.
5. Salam A., Strathdee J., On superfields and Fermi-Bose symmetry, Preprint IC/74/42, Trieste, 1974.
6. Березин Ф. А., Кац Г. И., Матем. сб., **82(124)**, 343 (1970).
7. Corvin L., Ne'eman Y., Sternberg S., Graded Lie algebras in mathematics and physics (Bose-Fermi symmetry), Preprint TAUP-448-74, Tel Aviv, 1974.
8. Ainsaar A., A supersymmetry of bosons, Preprint FI-38, Tartu, 1975.

Academy of Sciences of the Estonian SSR
Institute of Physics

Received
Apr. 30, 1975

E. Vilde Pedagogical Institute, Tallinn