

<https://doi.org/10.3176/phys.math.1978.3.01>

O. VAARMANN

УДК 517.948 : 513.88.518

CERTAIN ITERATIVE METHODS FOR THE NONLINEAR LEAST SQUARES PROBLEM

This paper is concerned with the problem of solving nonlinear operator equation in the sense of least squares.

Let $F(x)$ be a differentiable (sufficiently often with respect to x) operator of one Hilbert space, H_1 , another, H_2 , and such that inverse $[F'(x)]^{-1}$ may fail to exist.

To solve the equation

$$[F'(x)]^* F(x) = 0 \quad (1)$$

we consider the iterative methods of type

$$x_{k+1} = x_k - Q(x_k, A_k) \quad (k=0, 1, \dots), \quad (2)$$

where $Q(x, A_k)$ — is an operator of H_1 into itself and A_k — is an approximation to the pseudoinverse $[F'(x_k)]^+$ with

$$A_k = A_k P_h, \quad \|(P_h - F'(x_k) A_k) F(x_k)\| \leq \gamma_k \|P_h F(x_k)\| \quad (\gamma_k < 1). \quad (3)$$

Without proof we shall state a basic convergence theorem concerning the methods of type (2) examined in [1-2]. We shall construct two rapidly convergent iterative methods like those of [1-5] not requiring linear operators to be inverted and consider also some acceleration possibilities discussed in [6] (see methods 4° and 6° in section 3). Making use of the basic theorem we shall prove their convergence to a solution of the equation (1), establish their convergence rates and illustrate the methods by some numerical examples.

1. Basic theorem. For a closed subspace S of a Hilbert space H , we let by P_S denote the orthogonal projector of H onto S . Throughout this paper we shall assume that the range $R(F'(x))$ is closed; for simplicity we shall use the following notations $R(x) = R(F'(x))$, $R^*(x) = R([F'(x)]^*)$, $P_h = P_{R(F'(x_k))}$ and $P_h^* = P_{R([F'(x_k)]^*)}$.

Assume that $\{x_k\}$ generated by (2) satisfy the condition

$$\|P_{h+1} F(x_{k+1})\| \leq \sum_{v=1}^{n_1} \omega_v \gamma^{a_v} \|P_h F(x_k)\|^{b_v} + \sum_{i=p}^{n_2} c_i \|P_h F(x_k)\|^i \quad (a_v, b_v \geq 0), \quad (A)$$

and the quantities γ_k can be expressed in the form of

$$\gamma_{k+1} = (\gamma_k + h \|P_h F(x_k)\|)^q, \quad \gamma_0 + h \|P_0 F(x_0)\| < 1, \quad (B)$$

or vanish in the limit as $k \rightarrow \infty$ according to another rule, or $\gamma_k \leq \gamma < 1$

($k=0, 1, \dots$) while γ can as well amount to zero. There ω_v, c_i, h are certain nonnegative constants with the exception of $c_p > 0$.

If the quantities γ_h are defined by (B), we shall introduce sequences $\{b_h\}$ and $\{d_h\}$. Let be $s = \min_v b_v$ and $l = \min_{v=s} a_v$ for all terms in (A) with

$\omega_v \neq 0$. We define for $a_v + b_v \geq p$ with $\omega_v \neq 0$ and $q \geq p$

$$b_h \geq \max\{\gamma_h, \|P_h F(x_h)\|\}, \quad (4)$$

$$d_h = \max\left\{\sum_{v=1}^{n_1} \omega_v b_h^{a_v + b_v - p} + \sum_{i=p}^{n_2} c_i b_h^{i-p}, (1+h)^q b_h^{q-p}\right\}, \quad (5)$$

and for $l < p - s$ and $q \geq \frac{p(p-s)}{l}$

$$b_h \geq \max\left\{\gamma_h^{p-s}, \|P_h F(x_h)\|\right\}, \quad (6)$$

$$d_h = \max\left\{\sum_{v=1}^{n_1} \omega_v b_h \frac{(p-s)a_v + l(b_v - p)}{l} + \sum_{i=p}^{n_2} c_i b_h^{i-p}, (1+h)^{\frac{ql}{p-s}} b_h^{\frac{ql}{p-s} - p}\right\}. \quad (7)$$

On the occasion of $\gamma_h = 0$, we denote by

$$d_h = c_p + \sum_{i=p+1}^{n_2} c_i \|P_h F(x_h)\|^{i-p} = c_p + \sum_{i=p+1}^{n_2} c_i d_h^{i-p}. \quad (8)$$

If only any coefficient $\omega_v \neq 0$, then in the case of $\gamma_0 \geq \gamma_1 \geq \dots \geq \gamma_h \geq \dots \geq 0$ we take

$$\delta_k^{(4)} = \sum_{v=1}^{n_1} \omega_v \gamma_h^{a_v} \|P_h F(x_h)\|^{b_v - 1} + \sum_{i=p}^{n_2} c_i \|P_h F(x_h)\|^{i-1}. \quad (9)$$

If there is at least a coefficient $\omega_v \neq 0$ with $b_v \geq 1$ and $\gamma_h \leq \gamma < 1$, then γ stands for γ_h in (9).

In this paper we shall use for the following a theorem that is not hard to prove [1].

Theorem 1. Let $x_0 \in H_1$, $S = \{x \in H_1 : \|x - x_0\| \leq \varrho\}$ and let $P_{R(x)} F(x)$ be a continuous operator fulfilling condition (A) and the inequality

$$\|Q(x_h, A_h)\| \leq \lambda \|P_h F(x_h)\| \quad (0 < \lambda < \infty)$$

be valid in S .

Then 1) If $p=1$ or $\gamma_h \leq \gamma \neq 0$, $r_1 = \lambda \|P_0 F(x_0)\| / (1 - \delta) \leq \varrho$, $\delta = \delta_0^{(4)} < 1$ then the equation $[F'(x)]^* F(x) = 0$ has a solution x^* in S , to which the sequence (2) converges with $\|x^* - x_0\| \leq r_1$ and

$$\|x_h - x^*\| \leq r_1 \delta^h;$$

if $p \geq 2$ provided $\gamma_0 \geq \gamma_1 \geq \dots \geq \gamma_h \geq \dots \geq 0$, then $\delta_k^{(4)} \searrow 0$ and the sequence (2) converges superlinearly, i. e.,

$$\|x_h - x^*\| \leq r_1 \prod_{i=0}^{h-1} \delta_i^{(4)};$$

2) if $p \geq 2$ and quantities γ_h ($k=0, 1, \dots$) are expressed in the form (B)

or $\gamma_k=0$, ($k=0, 1, \dots$), $\delta = \frac{p-1}{\sqrt[p-1]{d_0 b_0}} < 1$, $r_p \leq \lambda H_0(\delta) / \sqrt[p-1]{d} \leq \rho$ where $d = \lim_{k \rightarrow \infty} d_k$ and

$$H_k(\delta) = \sum_{i=k}^{\infty} \delta^{p^i}$$

then the equation $[F'(x)]^*F(x) = 0$ has a solution x^* in S , to which the sequence (2) converges, with $\|x^* - x_0\| \leq r_p$ and

$$\|x_k - x^*\| \leq (\lambda \sqrt[p-1]{d}) H_k(\delta).$$

From this theorem, in a special case, $A_k = [F'(x_k)]^+$, the convergence for the analogues of the generalizations of the well-known rapidly convergent iterative methods, such as Newton's, Chebyshev's methods, etc., to a solution of the equation (1) concludes, and estimations for the rate of convergence follow [1, 2].

Corollary. Let τ be a real number, $2 \leq \tau < p - 1$. If a) $q \geq \tau$ and $a_\nu + b_\nu \geq \tau$ with $\omega_\nu \neq 0$ or b) $q \geq \frac{\tau(\tau - s)}{l}$ and $l < \tau - s$, then the order of the convergence for iterative procedure (2) is equal to τ .

2. We shall now state two facts which will be useful in the sequel.

Remark 1. Suppose that $F'(x)$ and $[F'(x)]^+$ satisfy Lipschitz condition in a certain region S :

$$\|F'(x) - F'(y)\| \leq L_1 \|x - y\| \quad (x, y \in S), \tag{10}$$

$$\|[F'(u)]^+ - [F'(v)]^+\| \leq L \|u - v\| \quad (u, v \in S), \tag{11}$$

then

$$\|P_{R(x)} - P_{R(y)}\| \leq L_0 \|x - y\|, \tag{12}$$

$$\|(P_{h+1} - P_{h+1}P_h)F(x_h)\| \leq G \|P_h F(x_h)\|^2, \tag{13}$$

where $G = \lambda L_0(1 + \lambda L_0 F)$ and $F = \sup_{x \in S} \|F(x)\|$.

Remark 2. If $F(x)$ is Lipschitz continuous and

$$\|(P_{R(y)} - P_{R(y)}P_{R(x)})F(x)\| \leq N_0 \|x - y\|, \tag{14}$$

then $P_{R(x)}F(x)$ is continuous [1].

If (10) and (11) are valid, then the continuity of $P_{R(x)}F(x)$ follows from (12).

For solving (1) we now make use of the following iterative method:

$$x_{k+1} = x_k - D_k [F'(x_k)]^* F(x_k), \tag{15}$$

$$D_{k+1} = 2D_k - D_k [F'(x_{k+1})]^* F'(x_{k+1}) D_k \tag{16}$$

where the starting operator D_0 is close enough to $\{[F'(x_0)]^* F'(x_0)\}^+$ with $D_0 = P_0^* D_0 = D_0 P_0^*$ [7].

Since $F'(x)$ has a closed range, then it has the bounded pseudo-inverse $[F'(x)]^+$ [8]. Let β_k ($k=0, 1, \dots$) C and N be quantities satisfying the inequalities

$$\|P_{k+1} - F'(x_{k+1})A_k\| \leq \beta_k, \quad \|[F'(x)]^+\| \leq C, \quad x \in S, \tag{17}$$

$$\|P_{R(y)} - P_{R(y)}P_{R(x)}\| \leq N, \quad \|P_{R^*(y)} - P_{R^*(y)}P_{R^*(x)}\| \leq N, \quad (x, y \in S). \quad (18)$$

Denote by

$$\gamma_0 = \max\{(\beta_0 + h_1 \|P_0 F(x_0)\|)^2, \|P_0 - F'(x_0)A_0\|\}, \quad b_h = \max\{\gamma_h, \|P_h F(x_h)\|\}. \quad (19)$$

Expressions for γ_h and β_h will be given below.

Putting

$$A_h = D_h [F'(x_h)]^*, \quad B_h = [F'(x_h)]^* F'(x_h) \quad (20)$$

as a corollary of Theorem 1 it follows

Theorem 2. Let $x_0 \in H_1$, $S = \{x \in H_1 : \|x - x_0\| \leq \rho\}$ and let in S be valid the following conditions:

1° operator $F(x)$ has Fréchet-derivative $F'(x)$;

2° derivative $F'(x)$ satisfies Lipschitz condition, i. e.,

$$\|F'(x) - F'(y)\| \leq L_1 \|x - y\| \text{ for all } x, y \in S.$$

1) If the operators $F'(x)$ and $F'(y)$ are in the acute case* and for any $x, y \in S$ $\|(P_{R(y)} - P_{R(y)}P_{R(x)})F(x)\| \leq N_0 \|x - y\|$, $\beta_1 = \gamma_0 + N + L_1 \lambda^2 \|P_0 F(x_0)\| \leq$

$$\leq \beta_0, \quad \delta = \delta_0^{(1)} = \gamma_0 + N_0 \lambda + \frac{1}{2} L_1 \lambda^2 \|P_0 F(x_0)\| < 1, \quad r_1 = \lambda \|P_0 F(x_0)\| / (1 - \delta) \leq \rho,$$

then the equation $[F'(x)]^* F(x) = 0$ has in S a solution x^* , to which the sequence (15)–(16) converges with $\|x^* - x_0\| \leq r_1$ and

$$\|x_k - x^*\| \leq r_1 \delta^k;$$

2) If $\|[F'(x)]^+ - [F'(y)]^+\| \leq L \|x - y\|$, $\beta_1 = \gamma_0 + h_1 \|P_0 F(x_0)\| \leq \beta_0$, $L_0 r_2 < 1$,

$$\delta = \delta^{(2)} = db_0 < 1, \quad r_2 = \lambda H_0(\delta) / d, \text{ where } h_1 = L_0 \lambda + L_1 \lambda^2 \text{ and } H_h(\delta) = \sum_{i=h}^{\infty} \delta^{2^i},$$

then the sequence (15)–(16) converges quadratically with $\|x^* - x_0\| \leq r_2$ and

$$\|x_k - x^*\| \leq r_2 H_h(\delta) / d,$$

where $d = \max\{(1 + h_0 + h_1)^2, 1 + h_2\}$, $h_0 = C(1 + \beta_0) \lambda L_1 / (1 - L_2 r_2)$, $h_2 = G + \frac{1}{2} L_1 \lambda^2$. If, in addition, to Lipschitz continuity of $[F'(x)]^+$ the

inequalities $R(x_k) \supseteq R(x_{k+1})$ are fulfilled, then $h_1 = L_1 \lambda^2$ and $h_2 = \frac{1}{2} L_1 \lambda^2$.

Proof. Let T be a linear bounded operator from H_1 into H_2 with the closed range $R(T)$.

It is well-known that

$$T^{*+} = T^{*+}, \quad P_{R(T)} = TT^+ = T^{*+} T^*, \quad T^+ = T + P_{R(T)}, \quad T^* = T^* P_{R(T)}, \\ P_{R(T^*)} T^* = T^*, \quad P_{R(T^*)} = T^+ T. \quad (21)$$

From (16) and (20) it is evident that $P_0^* A_h = A_h$ provided $D_0 = P_0^* D_0$. Apparently, then $P_h^* A_h = P_h^* P_0^* A_h$ and by (3), (20)–(21) we have

$$\|Q(x_k, A_h)\| = \|A_h\| = \|[F'(x_k)]^+ [P_h + F'(x_k)A_h - P_h + (P_0^* - P_0^* P_h^*) A_h]\|. \quad (22)$$

Moreover by (16), (17), (20)

* Definition. The linear operators A and B are said to be in the acute case if $R(A)$ and $R(B)$ are in the acute case as well as $R(A^*)$ and $R(B^*)$ [9].

$$\begin{aligned} \|F'(x_h)D_{h-1}P_{h-1}^*\| &= \|F'(x_h)A_{h-1}[F'(x_{h-1})]^{*+}\| \leq C(1+\beta_{h-1}), \\ \|F'(x_h)D_{h-1}\| &= \|F'(x_h)D_{h-1}P_0^*\| = \|F'(x_h)D_{h-1}P_{h-1}^* + \\ &+ F'(x_h)D_{h-1}(P_0^* - P_0^*P_{h-1}^*)\|. \end{aligned} \tag{23}$$

On the basis of Taylor expression

$$\begin{aligned} P_{h+1}F(x_{h+1}) &= (P_{h+1} - P_{h+1}P_h)F(x_h) + P_{h+1}\{(P_h - F'(x_h)A_h)F(x_h) + \\ &+ \int_0^1 [F'(x_h) - F'(x_h + t(x_{h+1} - x_h))]A_hF(x_h) dt\}. \end{aligned} \tag{24}$$

Using (16)–(21), we obtain

$$\begin{aligned} \|P_h - F'(x_h)A_h\| &= \|P_h - F'(x_h)D_h[F'(x_h)]^*\| \leq \\ &\leq \|(P_h - F'(x_h)D_{h-1}[F'(x_h)]^*)\|^2, \end{aligned} \tag{25}$$

$$\begin{aligned} \|P_h - F'(x_h)D_{h-1}[F'(x_h)]^*\| &\leq \|P_h - F'(x_h)A_{h-1}\| + \\ &+ \|F'(x_h)D_{h-1}([F'(x_{h-1})]^* - [F'(x_h)]^*)\|. \end{aligned} \tag{26}$$

It is plain that we may write

$$\begin{aligned} \|P_h - F'(x_h)A_{h-1}\| &\leq \|P_h(P_{h-1} - F'(x_{h-1})A_{h-1})\| + \|(P_h - P_hP_{h-1})\| + \\ &+ \|F'(x_h) - F'(x_{h-1})\| \|A_hF(x_h)\|. \end{aligned} \tag{27}$$

1) If $F'(x)$ and $F'(y)$ are in the acute case, then $N < 1$ [9] and this with (22), (23), (27) and $\delta < 1$ implies that

$$\begin{aligned} \|A_k\| &\leq C(1+\gamma_h)/(1-N), \quad \|F'(x_h)D_{h-1}\| \leq C(1+\beta_{h-1})/(1-N), \\ \|P_h - F'(x_h)A_{h-1}\| &\leq \gamma_{h-1} + N + \lambda^2L_1\|P_{h-1}F(x_{h-1})\|. \end{aligned}$$

Therefore it is sufficient to take for $\lambda = \sup\{\lambda_h\}$ with $\lambda_h = C(1+\gamma_h)/(1-N)$ and

$$\beta_{h-1} = \gamma_{h-1} + N + \lambda^2L_1\|P_{h-1}F(x_{h-1})\|.$$

Because of (25), (26) and $\beta_h \leq \beta_0$ we get

$$\|P_h - F'(x_h)D_{h-1}[F'(x_h)]^*\| \leq \beta_{h-1} + h_0\|P_{h-1}F(x_{h-1})\|,$$

where $h_0 = C(1+\beta_0)\lambda L_1/(1-N)$, and we can determine γ_h by $\gamma_h = (\beta_{h-1} + h_0\|P_{h-1}F(x_{h-1})\|)^2$. From (24) we see that

$$\|P_{h+1}F(x_{h+1})\| \leq (\gamma_h + N_0\lambda + \frac{1}{2}L_1\lambda^2\|P_hF(x_h)\|)\|P_hF(x_h)\| = \delta_h\|P_hF(x_h)\|.$$

Since $\delta_0 < 1$, $\beta_1 \leq \beta_0$ we obtain by induction that $\beta_h \leq \beta_0$, $\gamma_h \leq \gamma_0$, $\lambda_h \leq \lambda_0 = C(1+\gamma_0)/(1-N) = \lambda$ and $\delta_h \leq \delta_0 < 1$.

2) Now taking into account Remark 1, we have

$$\begin{aligned} \|P_h - P_hP_{h-1}\| &\leq \|P_h - P_{h-1}\| \leq L_0\lambda\|P_{h-1}F(x_{h-1})\|, \\ \|P_0 - P_h\| &\leq L_0\|x_0 - x_h\| \leq L_0r_2, \end{aligned}$$

so that

$$\beta_{h-1} = \gamma_{h-1} + h_1\|P_{h-1}F(x_{h-1})\|, \quad \lambda = C(1+\gamma_0)/(1-L_0r_2),$$

$$\|P_{h+1}F(x_{h+1})\| \leq \gamma_h\|P_hF(x_h)\| + \left(G + \frac{1}{2}L_1\lambda^2\right)\|P_hF(x_h)\| \leq (1+h_2)b_h^2, \tag{28}$$

where $h_1 = L_0\lambda + L_1\lambda^2$ and $h_2 = G + \frac{1}{2}L_1\lambda^2$.

Since

$$\|(P_h - F'(x_h)A_h)\| \leq (\beta_{h-1} + h_0\|P_{h-1}F(x_{h-1})\|)^2$$

with $h_0 = C(1 + \beta_0)\lambda L_1 / (1 - L_0r_2)$, we can define γ_h by

$$\gamma_h = (\gamma_{h-1} + h\|P_{h-1}F(x_{h-1})\|)^2, \quad h = h_1 + h_0.$$

From (24) and (27) it is evident that $s = l = 1$ and $p = 2$.

With a consideration of all these results, a straightforward application of Theorem 1 yields Theorem 2.

Consider

$$x_{h+1} = x_h - A_h F(x_h), \quad (29)$$

$$D_{h+1} = D_h + \alpha_{h+1}(I_1 - B_{h+1}D_h), \quad (30)$$

where A_h and B_{h+1} are as in (20), I_1 — identity operator in H_1 , α_{h+1} — is a real number, $0 < \alpha_{h+1} < 2/M^2$, $\|F'(x)\| \leq M$.

Denote

$$\gamma_0 = \max\{\mu\beta_0 + \mu h_1\|P_0F(x_0)\|, \|P_0 - F'(x_0)A_0\|\}, \quad (31)$$

where $\mu = \sup_h \{\mu_h\}$, $\|P_h - \alpha_h F'(x_h)[F'(x_h)]^*\| \leq \mu_h$.

Theorem 3. Let $x_0 \in H_1$, $S = \{x \in H_1 : \|x - x_0\| \leq \varrho\}$, and let in S the conditions 1° — 2° of Theorem 2 be valid.

If $\|[F'(x)]^+ - [F'(y)]^+\| \leq L\|x - y\|$, $R^*(x_0) \supseteq R^*(x)$, $\beta_1 = \gamma_0 + h_1\|P_0F(x_0)\| \leq \beta_0$, $\delta = \gamma_0 + h_2\|P_0F(x_0)\| < 1$, $L_0r < 1$, $r = \lambda\|P_0F(x_0)\| / (1 - \delta) \leq \varrho$, then the equation $[F'(x)]^*F(x) = 0$ has, in S , a solution x^* , to which the sequence (29) — (30) converges superlinearly with $\|x^* - x_0\| \leq r_1$, and

$$\|x_k - x^*\| \leq r_1 \prod_{i=0}^{k-1} \delta_i,$$

where $\delta_i = \gamma_i + h_2\|P_iF(x_i)\|$, and h_i ($i = 1, 2$) are as in Theorem 2.

Proof. From (29) — (31) it follows easily that

$$\begin{aligned} \|P_h - F'(x_h)A_h\| &= \|(P_h - \alpha_h F'(x_h)[F'(x_h)]^*) \times \\ &\times (P_h - F'(x_h)D_{h-1}([F'(x_{h-1})]^* + [F'(x_h)]^* - [F'(x_{h-1})]^*))\| \leq \\ &\leq \mu(\beta_{h-1} + h_0\|P_{h-1}F(x_{h-1})\|). \end{aligned}$$

The condition $R^*(x_0) \supseteq R^*(x)$ is only needed for bounding $\|A_h\|$, because then $P_0^*D_h = D_h$.

The rest of it is analogous to proof of Theorem 2.

3. Note that for choosing $\{A_h\}$ we can employ iterative procedures of the higher order $q \geq 2$, and for defining $Q(x, A_h)$ we can make use of a formula generating iterative process of the order $p \geq 2$ following [1-5]. Because of their susceptibility to errors, the convergence rate of iterative methods of a higher order is probably too slow to be of practical value in the actual solution, considering the labour involved in obtaining the iterates.

As applied to the solution of nonlinear systems of an algebraic system, the methods (15) — (16) and (29) — (30) require only slightly higher computational efforts than those of [10]. However, we proved

the convergence of methods (15)–(16) and (29)–(30) to a solution of the equation (1) under weaker assumptions and achieved better bounds on the rate of convergence than by methods of [1].

Finally, note that it is difficult to say which is the best criterion for comparing certain computational techniques, and that is why in this paper no claim is made for judging the efficiency of methods.

As an illustration of the above methods, we consider some computational experiments. The methods were tested on the following problems taken from the literature [14].

Example 1. The system of equations is

$$F(x) = \begin{cases} f_1 = x_1^2 + x_2^2 - 2 = 0, \\ f_2 = x_1 - x_2 = 0, \\ f_3 = x_1 x_2 - 1 = 0, \end{cases}$$

whose solutions are (1, 1) and (-1, -1).

Example 2.

$$F(x) = \begin{cases} \tilde{f}_1 = x_1^2 + x_2^2 - 2 = 0, \\ \tilde{f}_2 = (x_1 - 2)^2 + x_2^2 - 2 = 0, \\ \tilde{f}_3 = (x_1 - 1)^2 + x_2^2 - 9 = 0, \end{cases}$$

whose least squares solutions are (1.00000; ±1.914854) and $\min \sum_i^3 \tilde{f}_i^2 = 42.666\ 667$.

For solving this equations, we apply the following algorithms

- 1° $x_{h+1} = x_h - [F'(x_h)]^+ F(x_h)$, 5° $x_{h+1} = x_h - D_h [F'(x_h)]^* F(x_h)$,
- 2° $x_{h+1} = x_h - [F'(x_0)]^+ F(x_h)$, $D_{h+1} = 2D_h - D_h B_{h+1} D_h$,
- 3° $x_{h+1} = x_h - D_h [F'(x_h)]^* F(x_h)$, 6° $x_{h+1} = x_h - [2D_h - D_h B_h D_h] [F'(x_h)]^* F(x_h)$,
 $D_{h+1} = D_h + \alpha_{h+1} (I_1 - B_{h+1} D_h)$, $D_{h+1} = 2D_h - D_h B_{h+1} D_h$,
- 4° $x_{h+1} = x_h - [2D_h - D_h B_h D_h] [F'(x_h)]^* F(x_h)$,
 $D_{h+1} = D_h + \alpha_{h+1} (I_1 - B_{h+1} D_h)$,

where $\alpha_h = \frac{3}{2M_h}$, $M_h = \max_i \sum |b_{ij}^{(h)}|$, $[F'(x_h)]^* F'(x_h) = (b_{ij}^{(h)})$.

The algorithms were considered to have converged to the desired solution when the inequality $\|x_{h+1} - x_h\| \leq \varepsilon = 0.000\ 0010$ was satisfied. In the table we give the number of iterations required to obtain convergence.

Example	Initial guess	1°	2°	3°a	3°b	4°a	4°b	5°a	5°b	6°a	6°b
1	3	6	26	9	11	7	9	8	10	7	9
2	2										
2	10	8	95	28	13	(-)	12	(-)	15	(-)	14
2	20										
2	1,5	5	7	8	10	6	8	7	8	6	6
	2										

The figures *a* and *b* denote that the process was started with $D_0 = B_0^+$ and $D_0 = \alpha_0 I_1$ respectively, but (-) fails to converge.

REFERENCES

1. Ваарманн О. О сходимости методов с последовательной аппроксимацией псевдообратного оператора. № 3094-76 Деп. Новосибирск, 1976.
2. Ваарманн О. О сходимости итерационных методов с использованием псевдо-

- обратного оператора или его аппроксимации. — В кн.: Тезисы II Всесоюз. семинара, 31 мая — 3 июня 1976, Харьков. Харьков, 1976, с. 95—99.
3. Лика Д. К., Шафиев Р. А. Об усиленном методе Ньютона—Канторовича с последовательной аппроксимацией в гильбертовом пространстве. — В кн.: Приближенное решение уравнений. Кишинев, 1973, с. 24—29.
 4. Шафиев Р. А. Об аппроксимационном аналоге усиленного метода Ньютона—Канторовича. — Изв. АН МолдССР, Сер. физ.-техн. и матем., 1975, № 3, с. 16—23.
 5. Vaarmann, O. On methods with successive approximation of the pseudoinverse operator. — In: Colloquia Mathematica Societatis János Bolyai (12. Progress in Operations Research). Eger (Hungary), 1974, p. 941—946.
 6. Ваарманн О. Об ускорении методов с последовательной аппроксимацией псевдообратного оператора. — Изв. АН ЭССР, Физ. Матем., 1975, т. 24, № 2, с. 239—242.
 7. Showalter, D. W., Ben-Israel, A. Representation and computation of the generalized inverse of a bounded linear operator between two Hilbert spaces. — Atti Accad. Naz. Lincei. Mem., Ser. 8, 1970, v. 48, p. 184—194.
 8. Desoer, C. A., Whalen, B. H. A note on pseudoinverses. — SIAM J. Appl. Math., 1963, v. 11, N 2, p. 442—447.
 9. Wedin, P.-A. Perturbation theory for pseudoinverses. — BIT, 1973, v. 13, p. 217—232.
 10. Ваарманн О. О применении обобщенных обратных операторов и их приближений для решения нелинейных уравнений. — Изв. АН ЭССР, Физ. Матем., 1970, т. 19, № 3, с. 265—274.
 11. Ben-Israel, A. A Newton—Raphson method for the solution of systems of equations. — J. Math. Anal. Appl., 1966, v. 15, N 2, p. 243—253.

*Academy of Sciences of the Estonian SSR,
Institute of Cybernetics*

Received
Apr. 25, 1977

O. VAARMANN

MÕNED ITERATSIOONIMEETODID MITTELINEAARSE VÄHIMRUUTUDE ÜLESANDE LAHENDAMISEKS

Mittelineaarse vähimruutude ülesande lahendamiseks Hilberti ruumis on konstrueeritud kaks iteratsioonimeetodit (koonduvusjärk $p > 1$), mis põhinevad pseudopöördoperaatori järkjärgulisel aproksimeerimisel. On tõestatud valemitega (15)—(16) ja (29)—(30) defineeritud iteratsiooniprotsesside koonduvus võrrandi (1) lahendiks ning antud koondumiskiruse hinnangud. Teoreetiliste väidete illustreerimiseks on lahendatud kaks näiteülesannet, mis käsitlevad mittelineaarsete võrrandisüsteemide lahendi määramist vähimruutude mõttes.

O. ВААРМАНН

НЕКОТОРЫЕ ИТЕРАЦИОННЫЕ МЕТОДЫ ДЛЯ РЕШЕНИЯ НЕЛИНЕЙНОЙ ЗАДАЧИ НАИМЕНЬШИХ КВАДРАТОВ

Для решения нелинейной задачи наименьших квадратов построены два быстроходящихся итерационных метода с последовательной аппроксимацией псевдообратного оператора. Доказана сходимость итерационных процессов (15)—(16) и (29)—(30) к решению уравнения (1) и установлены оценки для их скорости сходимости. Теоретические соображения иллюстрированы числовыми примерами.