

Interpolation of approximation spaces with nonlinear projectors

Irina Asekritova

Department of Mathematics, Växjö University, 351 95 Växjö, Sweden; ias@msi.vxu.se

Received 10 April 2006

Abstract. Approximation spaces defined by multiparametric approximation families with possible nonlinear projectors are considered. It is shown that a real interpolation space for a tuple of such spaces is again an approximation space of the same type.

Key words: interpolation functor, approximation space, K -functional.

Let $\vec{X} = (X_0, X_1, \dots, X_n)$ be a tuple of Banach (or quasi-Banach) spaces, i.e. each space X_i , $i = 0, 1, \dots, n$, is a Banach (or quasi-Banach) space linearly and continuously embedded in some linear topological space \mathcal{X} . As usual, the interpolation space $K_{\vec{\theta}, q}(\vec{X})$ is defined by the norm

$$\|x\|_{\vec{\theta}, q} = \left(\int_{\mathbb{R}_+^n} (t_1^{-\theta_1} \dots t_n^{-\theta_n} K(\vec{t}, x, \vec{X}))^q \frac{dt_1}{t_1} \dots \frac{dt_n}{t_n} \right)^{1/q},$$

where $\vec{\theta} = (\theta_0, \theta_1, \dots, \theta_n)$, $0 < \theta_i < 1$, $\theta_0 + \theta_1 + \dots + \theta_n = 1$, $\vec{t} = (t_1, \dots, t_n) \in \mathbb{R}_+^n$ and

$$K(\vec{t}, x, \vec{X}) = \inf_{x=x_0+\dots+x_n} (\|x_0\|_{X_0} + t_1 \|x_1\|_{X_1} + \dots + t_n \|x_n\|_{X_n})$$

is the K -functional of the tuple \vec{X} .

Let $X \subset \mathcal{X}$ be a Banach space and $\mathcal{A} = \{A_{\vec{m}} \subset X, \vec{m} \in \mathbb{Z}_+^d\}$ be a family of linear subspaces $A_{\vec{m}}$, where $\vec{m} = (m_1, \dots, m_d)$ is a d -dimensional index with non-negative coordinates $m_i \geq 0$. We assume that the index set is ordered in coordinatewise order, i.e. $\vec{m} \leq \vec{l}$ means that $m_i \leq l_i$ for $1 \leq i \leq d$.

Definition 1. We will say that (X, \mathcal{A}) is a d -parametric approximation family if $\{0\} = A_{\vec{0}} \subset A_{\vec{m}} \subset A_{\vec{l}}$ for $\vec{m} \leq \vec{l}$.

As usual, the approximation number $e_{\vec{k}}(x, X)$ for $x \in X$ is defined by the formula

$$e_{\vec{k}}(x, X) = \inf \{ \|x - a\|_X, a \in A_{\vec{k}} \cap X \}.$$

Let Φ be an ideal Banach space of functions $f : \mathbb{Z}_+^d \rightarrow \mathbb{R}$ such that

$$l_0(\mathbb{Z}_+^d) \subset \Phi \subset l_\infty(\mathbb{Z}_+^d),$$

where $l_0(\mathbb{Z}_+^d)$ is a space of functions with finite support.

Definition 2. The approximation space $E_\Phi(X, \mathcal{A})$ is defined by the norm

$$\|x\|_{E_\Phi(X, \mathcal{A})} = \left\| \{e_{\vec{k}}(x, X)\}_{\vec{k} \in \mathbb{Z}_+^d} \right\|_\Phi.$$

Note that one-parametric approximation spaces have been considered by many authors (see, e.g., [1–5]). In the paper [6] multiparametric approximation spaces were considered, and conditions (on an interpolation functor \mathcal{F} and approximation family \mathcal{A}) were given under which the interpolation space of a tuple $E_{\vec{\Phi}}(\vec{X}, \mathcal{A}) = (E_{\Phi_0}(X_0, \mathcal{A}), \dots, E_{\Phi_n}(X_n, \mathcal{A}))$ is again the approximation space of the same type, i.e.,

$$\mathcal{F}[E_{\vec{\Phi}}(\vec{X}, \mathcal{A})] = E_{\mathcal{F}[\vec{\Phi}]}(\mathcal{F}[\vec{X}], \mathcal{A}). \quad (1)$$

A natural condition on the interpolation functor that arises here is the so-called splitting condition, namely

$$\mathcal{F}[\vec{\Phi}(\vec{X})] = \mathcal{F}[\vec{\Phi}](\mathcal{F}[(\vec{X})]), \quad (2)$$

where $\vec{\Phi}(\vec{X}) = (\Phi_0(X_0), \dots, \Phi_n(X_n))$ is a tuple of vector-valued spaces $\Phi_i(X_i)$.

It is known that the “splitting condition” is not always fulfilled. The case where \mathcal{F} is a functor of real interpolation $\mathcal{K}_{\theta, q}$ and $\vec{\Phi} = (l_{q_0}^{\vec{s}_0}, \dots, l_{q_n}^{\vec{s}_n})$ is studied in [7] and [8]; this case is important for applications.

In [6] it was shown that the formula (1) holds for an interpolation functor \mathcal{F} satisfying the “splitting condition” (2) and for a multiparametric approximation family \mathcal{A} with some family of linear projectors. But in some cases, for example, when considering quasi-Banach spaces (see [9]), it is useful to have an analogous result for approximation families with nonlinear projectors.

Let us have d one-parametric approximation families

$$\mathcal{A}^{(k)} = \left\{ A_m^{(k)} \subset X_0 + \dots + X_n, m \in \mathbb{Z}_+ \right\}, \quad k = 1, \dots, d,$$

and let us consider a special d -parametric approximation family

$$\mathcal{A} = \left\{ A_{\vec{m}} = A_{m_1}^{(1)} + \dots + A_{m_d}^{(d)} \right\}.$$

Definition 3. We will say that (\vec{X}, \mathcal{A}) is complemented if there exists a family of (possibly nonlinear) operators $P_m^{(k)} : X_0 + \dots + X_n \rightarrow A_m^{(k)}$ such that

1. $P_m^{(k)} x = x$ if $x \in A_m^{(k)}$,
2. $P_{m_0}^{(k_0)} P_{m_1}^{(k_1)} = P_{m_1}^{(k_1)} P_{m_0}^{(k_0)}$,
3. $\|P_m^{(k)} x\|_{X_j} \leq \gamma \|x\|_{X_j}$ with γ independent of m, k, j , and x .

To formulate our first result, let us consider operators $Q_{\vec{m}} : X_0 + \dots + X_n \rightarrow A_{\vec{m}}$ given by the formula

$$Q_{\vec{m}} = I - \prod_{i=1}^d (I - P_{m_i}^{(i)})$$

and let us also define operators

$$\Delta Q_{\vec{m}} = \prod_{i=1}^d (Q_{\vec{m}+e_i} - Q_{\vec{m}}),$$

where $e_i, 1 \leq i \leq d$, is the standard basis in \mathbb{R}^d . Let $\vec{\Phi} = (\Phi_0, \dots, \Phi_n)$ be a tuple of ideal spaces Φ_i with the Fatou property

$$\left\| \lim_{n \rightarrow \infty} f_n \right\|_{\Phi_i} \leq \underline{\lim}_{n \rightarrow \infty} \|f_n\|_{\Phi_i}$$

and such that the operator S is bounded in each Φ_i :

$$(Sf)(\vec{k}) = \sum_{\vec{l} \geq \vec{k}} f(\vec{l}), \vec{k} \in \mathbb{Z}_+^d.$$

Then the following theorem is true.

Theorem 4. Suppose that (\vec{X}, \mathcal{A}) is complemented, the operators $P_m^{(k)}$ are linear for $k \leq d - 1$ and the operators $P_m^{(d)}$ possess the following property: for any decomposition $x = x_0 + \dots + x_n$ ($x_j \in X_j$) there exists a decomposition $P_m^{(d)} x = y_0^m + \dots + y_n^m$ such that

$$\|x_j - y_j^m\|_{X_j} \leq \gamma e_m(x_j; \mathcal{A}, X_j),$$

where $\gamma > 0$ is some constant independent of x and m . Then

$$K(\cdot, x; E_{\vec{\Phi}}(\vec{X}, \mathcal{A})) \approx K(\cdot, \{\Delta Q_{\vec{m}} x\}_{\vec{m}}; \vec{\Phi}(\vec{X})).$$

The next theorem shows that spaces considered above are stable under real interpolation.

Theorem 5. Suppose that the tuples $\vec{\Phi}, \vec{X}$ are such that for the interpolation functor $K_{\vec{\theta}, q}$ the “splitting condition” is fulfilled. Then if the conditions of Theorem 1 hold, we have the equality

$$K_{\vec{\theta}, q}(E_{\vec{\Phi}}(\vec{X}, \mathcal{A})) = E_{K_{\vec{\theta}, q}(\vec{\Phi})}(K_{\vec{\theta}, q}(\vec{X}), \mathcal{A}).$$

REFERENCES

1. Brudnyi, Yu. and Krugljak, N. A family of approximation spaces. *Issled. Teor. Funkts. Mnogikh Veshchestv. Perem. (Yaroslavl)*, 1978, 2, 15–42 (in Russian).
2. Brudnyi, Yu. and Krugljak, N. *Interpolation Functors and Interpolation Spaces*, Vol. I. North Holland, Amsterdam, 1991.
3. Nilsson, P. Reiteration theorems for real interpolation and approximation spaces. *Ann. Mat. Pura Appl.*, 1982, **132**, 291–330.
4. Pietch, A. Approximation spaces. *J. Approx. Theory*, 1981, **32**, 115–138.
5. Peetre, J. and Sparr, G. Interpolation of normed Abelian groups. *Ann. Mat. Pura Appl.*, 1972, **92**, 217–262.
6. Asekritova, I. and Brudnyi, Yu. Interpolation of several multiparametric approximation spaces. *J. Approx. Theory*, 2004, **129**, 182–206.
7. Asekritova, I. and Krugljak, N. Real interpolation of vector-valued spaces in non-diagonal case. *Proc. Amer. Math. Soc.*, 2005, **133**, 1665–1675.
8. Asekritova, I., Krugljak, N. and Nikolova, L. Lizorkin–Freitag formula for several weighted spaces and vector-valued interpolation. *Studia Math.*, 2005, **170**, 173–201.
9. DeVore, R. and Popov, V. Interpolation of approximation spaces. In *Constructive Function Theory*. Bulgarian Academy of Sciences, Sofia, 1988, 110–119.

Aproksimatsiooniruumid mittelineaarsete projektoritega ja nende interpolatsioon

Irina Asekritova

On vaadeldud mitmest parameetrist sõltuvate parvede võimalike mittelineaarsete projektorite poolt defineeritud aproksimatsiooniruumide. On näidatud, et selliste ruumide iga reaalne interpolatsiooniruum moodustab jälle sama tüüpi aproksimatsiooniruumi.